







INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : C07D 307/80, 333/56, 235/12, 333/16, 417/06, 491/04, 409/12, 405/12, 471/04, A61K 31/34, 31/38, 31/415, 31/44 // (C07D 491/04, 307:00, 221:00) (C07D 471/04, 235:00, 221:00)</p>	A2	<p>(11) International Publication Number: WO 99/58518</p> <p>(43) International Publication Date: 18 November 1999 (18.11.99)</p>
<p>(21) International Application Number: PCT/US99/10201</p> <p>(22) International Filing Date: 10 May 1999 (10.05.99)</p> <p>(30) Priority Data: 09/076,205 12 May 1998 (12.05.98) US</p> <p>(71) Applicant: AMERICAN HOME PRODUCTS CORPORATION [US/US]; Five Giralda Farms, Madison, NJ 07940-0874 (US).</p> <p>(72) Inventors: MALAMAS, Michael, Sotirios; 2443 Oleander Circle, Jamison, PA 18929 (US). McDEVITT, Robert, Emmett; 232 Haworth Place, Somerset, NJ 08873 (US). ADEBAYO, Folake, Oluwemimo; 216 Dorchester Drive, Cranbury, NJ 08512 (US).</p> <p>(74) Agents: MILOWSKY, Arnold, S.; American Home Products Corporation, Patent Law Dept.-2B, One Campus Drive, Parsippany, NJ 07054 (US) et al.</p>		<p>(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>Without international search report and to be republished upon receipt of that report.</i></p>
<p>(54) Title: BIPHENYL OXO-ACETIC ACIDS USEFUL IN THE TREATMENT OF INSULIN RESISTANCE AND HYPERGLYCEMIA</p>		
<div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>(e)</p> </div> <div style="text-align: center;">  <p>(f)</p> </div> <div style="text-align: center;">  <p>(g)</p> </div> <div style="text-align: center;">  <p>(h)</p> </div> <div style="text-align: right;"> <p>(I)</p> </div> </div>		
<p>(57) Abstract</p> <p>This invention provides compounds of Formula (I) having the structure wherein A is (a) or (b); B is carbon or nitrogen; D is oxygen, sulfur, or nitrogen; E is carbon or nitrogen; Y is a bond, methylene, C(O), or CH(OH); R¹ is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-, or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms; Het is (c) (d); G is oxygen, sulfur or nitrogen; R² and R^{2a} are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl; R³ and R⁴ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro, amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, -NR⁷(CH₂)_mCO₂H, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3 heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms; R⁵ is hydrogen, alkyl of 1-6 carbon atoms, -CH(R⁸)R⁹, -C(CH₂)_nCO₂R¹⁰, -C(CH₃)₂CO₂R¹⁰, -CH(R⁸)(CH₂)_nCO₂R¹⁰, -CH(R⁸)C₆H₄CO₂R¹⁰; R⁶ is alkylene of 1 to 3 carbon atoms; R⁷ is hydrogen or alkyl of 1 to 6 carbon atoms; R⁸ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein the alkyl moiety contains 1-6 carbon atoms; Q is (e), (f), (g), or (h); W is oxygen, sulfur, or nitrogen; R⁹ is -CO₂R¹², -CONHR¹², tetrazole, -PO₃R¹²; R¹⁰ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of 7-15 carbon atoms; R¹² is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms; m = 1-3; n = 1-6; with the proviso that when R¹ is halogen, Y is a bond; or a pharmaceutically acceptable salt thereof, which are useful in treating metabolic disorders related to insulin resistance or hyperglycemia.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

- 1 -

BIPHENYL OXO-ACETIC ACIDS USEFUL IN THE TREATMENT OF INSULIN RESISTANCE AND HYPERGLYCEMIA

BACKGROUND OF THE INVENTION

5

The prevalence of insulin resistance in glucose intolerant subjects has long been recognized. Reaven et al (*American Journal of Medicine* 1976, 60, 80) used a continuous infusion of glucose and insulin (insulin/glucose clamp technique) and oral glucose tolerance tests to demonstrate that insulin resistance existed in a diverse group of nonobese, nonketotic subjects. These subjects ranged from borderline glucose tolerant to overt, fasting hyperglycemia. The diabetic groups in these studies included both insulin dependent (IDDM) and noninsulin dependent (NIDDM) subjects.

10

Coincident with sustained insulin resistance is the more easily determined hyperinsulinemia, which can be measured by accurate determination of circulating plasma insulin concentration in the plasma of subjects. Hyperinsulinemia can be present as a result of insulin resistance, such as is in obese and/or diabetic (NIDDM) subjects and/or glucose intolerant subjects, or in IDDM subjects, as a consequence of over injection of insulin compared with normal physiological release of the hormone by the endocrine pancreas.

15

The association of hyperinsulinemia with obesity and with ischemic diseases of the large blood vessels (e.g. atherosclerosis) has been well established by numerous experimental, clinical and epidemiological studies (summarized by Stout, *Metabolism* 1985, 34, 7, and in more detail by Pyorala et al, *Diabetes/Metabolism Reviews* 1987, 3, 463). Statistically significant plasma insulin elevations at 1 and 2 hours after oral glucose load correlates with an increased risk of coronary heart disease.

20

25

Since most of these studies actually excluded diabetic subjects, data relating the risk of atherosclerotic diseases to the diabetic condition are not as numerous, but point in the same direction as for nondiabetic subjects (Pyorala et al). However, the incidence of atherosclerotic diseases in morbidity and mortality statistics in the diabetic population exceeds that of the nondiabetic population (Pyorala et al; Jarrett *Diabetes/Metabolism Reviews* 1989, 5, 547; Harris et al, Mortality from diabetes, in *Diabetes in America* 1985).

30

The independent risk factors obesity and hypertension for atherosclerotic diseases are also associated with insulin resistance. Using a combination of insulin/glucose clamps, tracer glucose infusion and indirect calorimetry, it has been demonstrated that the insulin resistance of essential hypertension is located in peripheral

35

- 2 -

tissues (principally muscle) and correlates directly with the severity of hypertension (DeFronzo and Ferrannini, *Diabetes Care* 1991, 14, 173). In hypertension of the obese, insulin resistance generates hyperinsulinemia, which is recruited as a mechanism to limit further weight gain via thermogenesis, but insulin also increases renal sodium reabsorption and stimulates the sympathetic nervous system in kidneys, heart, and vasculature, creating hypertension.

It is now appreciated that insulin resistance is usually the result of a defect in the insulin receptor signaling system, at a site post binding of insulin to the receptor. Accumulated scientific evidence demonstrating insulin resistance in the major tissues which respond to insulin (muscle, liver, adipose), strongly suggests that a defect in insulin signal transduction resides at an early step in this cascade, specifically at the insulin receptor kinase activity, which appears to be diminished (reviewed by Haring, *Diabetologia* 1991, 34, 848).

Protein-tyrosine phosphatases (PTPases) play an important role in the regulation of phosphorylation of proteins. The interaction of insulin with its receptor leads to phosphorylation of certain tyrosine molecules within the receptor protein, thus activating the receptor kinase. PTPases dephosphorylate the activated insulin receptor, attenuating the tyrosine kinase activity. PTPases can also modulate post-receptor signaling by catalyzing the dephosphorylation of cellular substrates of the insulin receptor kinase. The enzymes that appear most likely to closely associate with the insulin receptor and therefore, most likely to regulate the insulin receptor kinase activity, include PTP1B, LAR, PTP α and SH-PTP2 (B. J. Goldstein, *J. Cellular Biochemistry* 1992, 48, 33; B. J. Goldstein, *Receptor* 1993, 3, 1-15; F. Ahmad and B. J. Goldstein *Biophys. Biophys. Acta* 1995, 1248, 57-69).

McGuire et al. (*Diabetes* 1991, 40, 939), demonstrated that nondiabetic glucose intolerant subjects possessed significantly elevated levels of PTPase activity in muscle tissue vs. normal subjects, and that insulin infusion failed to suppress PTPase activity as it did in insulin sensitive subjects.

Meyerovitch et al. (*J. Clinical Invest.* 1989, 84, 976) observed significantly increased PTPase activity in the livers of two rodent models of IDDM, the genetically diabetic BB rat, and the STZ-induced diabetic rat. Sredy et al. (*Metabolism*, 44, 1074, 1995) observed similar increased PTPase activity in the livers of obese, diabetic ob/ob mice, a genetic rodent model of NIDDM.

The compounds of this invention have been shown to inhibit PTPases derived from rat liver microsomes and human-derived recombinant PTPase-1B (hPTP-1B) *in vitro*. They are useful in the treatment of insulin resistance associated with obesity,


- 3 -

glucose intolerance, diabetes mellitus, hypertension and ischemic diseases of the large and small blood vessels.

DESCRIPTION OF THE INVENTION


5

This invention provides a compound of formula I having the structure



I

wherein




B is carbon or nitrogen;

D is oxygen, sulfur, or nitrogen;

E is carbon or nitrogen;

Y is a bond, methylene, C(O), or CH(OH);

- 15 R¹ is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-, or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

R² and R²ᵃ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl;

- 25 R³ and R⁴ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro,

- 4 -


amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, $-NR^7(CH_2)_mCO_2H$, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3 heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms;

R^5 is hydrogen, alkyl of 1-6 carbon atoms, $-CH(R^8)R^9$, $-C(CH_2)_nCO_2R^{10}$, $-C(CH_3)_2CO_2R^{10}$, $-CH(R^8)(CH_2)_nCO_2R^{10}$, $-CH(R^8)C_6H_4CO_2R^{10}$

R^6 is alkylene of 1 to 3 carbon atoms

R^7 is hydrogen or alkyl of 1 to 6 carbon atoms,

R^8 is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein the alkyl moiety contains 1-6 carbon atoms;



W is oxygen, sulfur, or nitrogen;

R^9 is $-CO_2R^{12}$, $-CONHR^{12}$, tetrazole, $-PO_3R^{12}$;

R^{10} is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of 7-15 carbon atoms;

R^{12} is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;

$m = 1-3$;

$n = 1-6$;

with the proviso that when R^1 is halogen, Y is a bond;

or a pharmaceutically acceptable salt thereof, which are useful in treating metabolic disorders related to insulin resistance or hyperglycemia.

25

Pharmaceutically acceptable salts can be formed from organic and inorganic acids, for example, acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic, malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric, sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic, camphorsulfonic, and similarly known acceptable acids when a compound of this invention contains a basic moiety. Salts may also be formed from organic and


30













- 5 -

inorganic bases, preferably alkali metal salts, for example, sodium, lithium, or potassium, when a compound of this invention contains a carboxylate or phenolic moiety, or similar moiety capable of forming base addition salts.

- 5 Alkyl includes both straight chain as well as branched moieties. Halogen means bromine, chlorine, fluorine, and iodine. It is preferred that the aryl portion of the aryl or aralkyl substituent is a phenyl, naphthyl or 1,4-benzodioxan-5-yl group; with phenyl being most preferred. The aryl moiety may be optionally mono-, di-, or tri- substituted with a substituent selected from the group consisting of alkyl of 1-6 carbon atoms, 10 alkoxy of 1-6 carbon atoms, trifluoromethyl, halogen, alkoxycarbonyl of 2-7 carbon atoms, alkylamino of 1-6 carbon atoms, and dialkylamino in which each of the alkyl groups is of 1-6 carbon atoms, nitro, cyano, $-\text{CO}_2\text{H}$, alkylcarbonyloxy of 2-7 carbon atoms, and alkylcarbonyl of 2-7 carbon atoms.
- 15 The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry in Formula I, the present invention includes such optical isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and 20 pharmaceutically acceptable salts thereof.


- 6 -

When A is , the following compounds of A are preferred:



- 7 -

Preferred compounds of this invention are those compounds of Formula I, in which:




wherein


5 B is carbon or nitrogen;

D is oxygen or sulfur;

or



or



10

Specifically preferred compounds of the present invention are set forth below:

(4'-benzofuran-3-yl-biphenyl-4-yloxy)-acetic acid

15

2-(4'-benzofuran-3-yl-biphenyl-4-yloxy)-3-phenyl-propionic acid

[4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid

20

2-[4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-ol

[4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid

25

2-[4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

- 2-[4'-(2-butyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 5 2-[4'-(2-benzoyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 10 (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
tromethamine salt
- 15 (2R)- 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-propionic acid
- (R)-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid
- (2R)-2-{4'-[2-(4-fluoro-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-
20 propionic acid
- (2R)-2-{4'-[2-(4-methoxy-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-
phenyl-propionic acid
- 25 [4'-(2-butyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid
- (2R)-2-{4'-[2-(hydroxy-phenyl-methyl)-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-
propionic acid
- 30 (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
methyl ester
- 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid
- 35 (2R)-2-{4'-[2-(3,4-dimethoxy-benzyl)-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-
phenyl-propionic acid

- (2R)-2-{4'-[2-(2,4-dimethoxy-benzyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid
- 5 (2R) 2-{4'-[2-(2,2-dimethyl-benzo[1,3]dioxol-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid
- (2R)-2-{4'-[2-(3,4-dihydroxy-benzyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid
- 10 (2R)-2-[4'-(2-benzyl-thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 3-phenyl-2-[4'-(2-thiazole-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid
- 15 (2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid sodium salt
- (2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid
- 20 (2S)-2-[4'-(2-benzyl-furo[2,3]pyridin-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-ol
- 25 4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-ol
- (2R)-2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 30 (2R)-2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid
- (2R)-2-[4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid
- 35

- 10 -

[(4,4''-dimethoxy-5'-(2-(phenylmethyl)benzo[b]thien-3-yl)phenyl)[1,1';3',1''-terphenyl]-2'-yl)oxy]-acetic acid

5 [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-4''-methoxy[1,1';3',1''-terphenyl]-4'-yl)oxy]-acetic acid

[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-acetic acid

10 [4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-acetic acid

(2S)-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-phenyl-butyric acid

15 4-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-butyric acid

N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-carboxy-propyl]-phthalamic acid

20 N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-methoxycarbonyl-propyl]-phthalamic acid

(2R)-2-[4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-chloro-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid

25 (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-phenyl-butyric acid

(2R)-2-[4'-(2-benzyl-benzo[b]-thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid

30 (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid

(2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid

35

- 11 -

- (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid
- 5 (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-methyl-pentanoic acid
- 2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-hexanoic acid
- 10 (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-butyric acid
- (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-octanoic acid
- 15 (2S)-2-[4-(2-benzyl-benzofuran-3-yl)-4''-methoxy-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid
- [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-3'',4''-dimethoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- 20 [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-3''-methoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- [[3,3-dimethoxy-5-[4-[2-(phenylmethyl)benzyl-benzo[b]-thiophen-3-yl)-[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- 25 [4-(2-benzyl-benzo[b]-thiophen-3-yl)-4''-methoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- [4-(2-benzyl-benzo[b]-thiophen-3-yl)-3'',4''-dimethoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- 30 [4-(2-benzyl-benzo[b]-thiophen-3-yl)-3'',4'',5''-trimethoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- [4-(2-benzyl-benzo[b]-thiophen-3-yl)-[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid
- 35 4-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxymethyl]-benzoic acid

- 12 -

(2S)-2-[4'-(1-benzyl-1H-benzimidazol-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

- 5 (2R)-2-[4'-(3-benzyl-3H-imidazol[4,5-b]pyridin-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt

(2R)-2-[4'-(2-benzyl-4,5-dimethyl-thiophen-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

10

(2R)-2-[4'-(2-benzoyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

15

(2R)-2-[4'-(2-benzyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

(2R)-2-[4'-(2-(hydroxy-phenyl-methyl)-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

20

(2S)-2-[4'-(2-benzyl-benzofuarn-3-yl)-3,5-dimethyl-biphenyl-4-yloxy]-3-phenyl-propionic acid

(2S)-2-[4'-(5-acetyl-thiophen-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt


25

or a pharmaceutically acceptable salt thereof.

- 30 The compounds of this invention were prepared according to the following schemes from commercially available starting materials or starting materials which can be prepared using literature procedures. These schemes show the preparation of representative compounds of this invention.

- 13 -

Scheme I




- In Scheme I, commercially available phenols (1; R^2 as defined above) were treated with 4-bromophenylacetyl bromide (2) in the presence of potassium carbonate to produce ketones (3). Compounds (3) were treated with polyphosphoric acid at high temperatures (150 C°) to afford benzofurans (4) [ref. J. Med. Chem. 1989, 32, 1700-1707]. Benzofurans (4) were coupled with aryl boronic acids of general structure (5; R^3, R^4 are alkyl, aryl, trifluoromethyl, substituted aryl, nitro, carbocyclic 5 to 7 carbon atoms rings or heterocyclic rings 5 to 7 atom rings with from 1 to 3 heteroatoms selected from oxygen, nitrogen, and sulfur) using the Suzuki protocol [ref. Syn. Comm. 1981, 11, 513-519] to produce biphenyls (6). The aryl boronic acids are either commercially available or can be prepared according to known methodology [ref. J. Org. Chem, 1984, 49, 5237-5243]. Biphenyls (6) were converted to biphenyls (8) using two different synthetic approaches. In the first synthetic approach, biphenyls (6) were treated first with $n\text{-BuLi}$ and either aldehydes R^1CHO or "Weinreb" amides $\text{R}^1\text{CON(CH}_3\text{)OCH}_3$ (R^1 is as defined above with the exception of halogen), and secondly with boron tribromide in dichloromethane to produce biphenyls (8). In the

- 14 -

second synthetic approach, biphenyls (6) first converted to biphenyls (7) with boron tribromide in dichloromethane [ref. J. Org. Chem. 1974, 39, 1427-1429], and then biphenyls (7) were converted to biphenyls (8) by using n-BuLi and either aldehydes R¹CHO or "Weinreb" amides R¹CON(CH₃)OCH₃. The required aldehydes and "Weinreb" amides are either commercially available or can be prepared according to known synthetic methodology [ref. Tet. Lett. 1993, 34, 6215-6218]. Biphenyls (8) were converted to biphenyls (9) by reduction of the ketones (Y = CO) using the Wolff-Kishner reduction (NH₂NH₂, KOH; ref. Org. Reactions, 1948, 4, 378), or reduction of the secondary hydroxy group (Y = CHOH) with sodium borohydride in trifluoroacetic acid [ref. Syn. Comm. 1990, 20, 487-493].

Scheme II




15

In Scheme II, the biphenyl boronic acid (11) was prepared from 10 according to known methodology [ref. J. Org. Chem. 1984, 49, 5237-5243] using n-BuLi to generate the aryllithium intermediate which was subsequently treated with triisopropyl borate. Boronic acid (11) was coupled with 3-bromobenzothiophenes (prepared according to reference J. Am. Chem. Soc. 1950, 72, 571-574) using the Suzuki protocol [ref. Syn. Comm. 1981, 11, 513-519] to produce biphenyls (12). Biphenyls (12) were converted to the biphenyls (13) in a similar manner as described in Scheme I for the conversion of biphenyls (6) to biphenyls (9).

- 15 -


Scheme III



- 5 In Scheme III, thienyl and furyl aldehydes (14) are either commercially available or can be prepared from substituted furans or thiophenes. Furans or thiophenes undergo metallation at the 2-position with alkyllithium reagents [J. Chem. Soc. Perkin I, 1977, 887] which upon treatment first with a formylating agent (i.e., dimethylformamide), and secondly bromination according to known methodology [ref. J. Am. Chem. Soc. 1950, 72, 571-574] to afford the required compounds (14).
- 10 Coupling of aldehydes (14) with 4, 4'-methoxy biphenyl boronic acid using the Suzuki protocol [ref. Syn. Comm. 1981, 11, 513-519] gave biphenyls (15). Known or easily prepared Grignard Reagents [ref. Chem. Re. 1954, 54, 835] $R^1Mg(Cl$ or $Br)$ (R^1 is as defined above with the exception of halogen, trifluoromethyl, lower alkoxy) were first
- 15 treated with aldehydes (14), followed by reduction of the produced methyl-hydroxy compounds with sodium borohydride and trifluoroacetic acid, and then demethylation with boron tribromide in dichloromethane to afford biphenyls (16).

- 16 -

Scheme IV




5 In Scheme IV, the furo[2,3-b]pyrines (18) (R^1 is aryl or alkyl) were prepared according to known methodology [ref. Tet. Lett. 1994, 35, 9355-9358]. Bromination of 18 with Br_2 in carbon tetrachloride produced 19. Pyridines (19) were converted to biphenyls (20) in a similar manner as described in Scheme I by Suzuki coupling with 4, 4'-methoxy biphenyl boronic acid, and demethylation with boron tribromide.

10

- 17 -


Scheme V



- 5 In Scheme V, the chloro-nitro-pyridines (21) were treated with primary amines to produce nitro-pyridines (22). The nitro-pyridines (22) were reduced with tin chloride to anilines (23), which upon treatment with ethyl chloroformate and sodium ethoxide gave imidazolones (24). The imidazolones (24) were converted to the 2-chloro-imidazole[4,5-b]pyridines (25). 2-Chlorobenzimidazoles (26) were alkylated in the
- 10 presence of sodium hydride to give benzimidazoles (27). Both, the imidazole[4,5-b]pyridines (25), and benzimidazoles (27) were converted to the biphenyls (28) in a similar manner as described in Scheme I by Suzuki coupling with 4, 4'-methoxy biphenyl boronic acid, and demethylation with boron tribromide.

- 18 -


Scheme VI



- 5 In Scheme VI, the biphenyl compounds (29; $Y = CO, CH_2$) can be monobrominated or dibrominated using bromine, potassium acetate and acetic acid. One equivalent of bromine in a high dilution reaction mixture and low temperatures in the range of 5-10 °C afforded predominantly the monobrominated product (30; $R^3, R^4 = H, Br$). The dibrominated product (30; $R^3, R^4 = Br, Br$) was obtained with two equivalents of bromine at room temperature. The Suzuki coupling protocol [ref. Syn. Comm. 1981, 11, 513-519] was used to generate the terphenyls 31 and 32. Coupling of the monobromo compounds (30; $R^3, R^4 = H, Br$) with boronic acids $R^{13}-Ar-B(OH)_2$; ($R^{13} = \text{halogen, trifluoromethyl, alkoxy, alkyl, nitro, amino, carboalkoxy}$) in the present of an inorganic base, for example $K_2CO_3, Ba(OH)_2$, and palladium (0 or II) catalyst, for example $Pd(PPh_3)_4, Pd(OAc)_2, (dppf)PdCl_2$, produced terphenyls (31; $R^3 = H$). Similarly, the dibromo compounds (30; $R^3, R^4 = Br, Br$) can undergo Suzuki coupling to afford either the di-coupled product (32) by using 2 equivalents of boronic acid at high temperatures (100 °C), or the mono-coupled-mono-bromo product (31; $R^3, R^4 = Br, Aryl-R^{13}$). Both the bromo and dibromo compounds can afford in the same synthetic manner products with various heterocyclic boronic acids, for example thiophene, furan, oxazole, thiazole, pyridine).

- 19 -

Scheme VII



5 In Scheme VII, the biphenyls (33) were alkylated with bromo or chloro-alkylcarboxylates [(Br or Cl)(CH₂)_nCO₂R¹²] in the presence of sodium hydride or potassium carbonate, using dimethylformamide or acetonitrile as the solvent. Subsequent saponification with sodium hydroxide in methyl alcohol and tetrahydrofuran produced biphenyls (34). Coupling of biphenyls (33) with hydroxy-

10 alkyl-carboxylates [HOCH(R⁸)CO₂R¹²] using the Mitsunobu protocol [ref. *Synthesis*. 1981, 1-27], followed by saponification with sodium hydroxide in methyl alcohol and tetrahydrofuran produced biphenyls (35). Similar experimental protocols were applied and for the biphenyls 16, 20, 28 and 32, for the production of compounds of Formula I.

15

The compounds of this invention are useful in treating metabolic disorders related to insulin resistance or hyperglycemia, typically associated with obesity or glucose intolerance. The compounds of this invention are therefore, particularly useful in the treatment or inhibition of type II diabetes. The compounds of this invention are

20 also useful in modulating glucose levels in disorders such as type I diabetes.

- 20 -

The ability of compounds of this invention to treat or inhibit disorders related to insulin resistance or hyperglycemia was established with representative compounds of this invention in the following two standard pharmacological test procedures which measure the inhibition of PTPase.

5

Inhibition of tri-phosphorylated insulin receptor dodecaphosphopeptide dephosphorylation by rat hepatic protein-tyrosine phosphatases (PTPases)

10 This standard pharmacological test procedure assess the inhibition of rat hepatic microsomal PTPase activity using, as substrate, the phosphotyrosyl dodecapeptide corresponding to the 1142-1153 insulin receptor kinase domain, phosphorylated on the 1146, 1150 and 1151 tyrosine residues. The procedure used and results obtained are briefly outlined below.

15 Preparation of Microsomal Fraction: Rats (Male Sprague-Dawley rats (Charles River, Kingston, NY) weighing 100-150 g, maintained on standard rodent chow (Purina)) are sacrificed by asphyxiation with CO₂ and bilateral thoracotomy. The liver is removed and washed in cold 0.85% (w/v) saline and weighed. The tissue is homogenized on ice in 10 volumes of Buffer A and the microsomes are isolated essentially as described by
20 Meyerovitch J, Rothenberg P, Shechter Y, Bonner-Weir S, Kahn CR. Vanadate normalizes hyperglycemia in two mouse models of non-insulin-dependent diabetes mellitus. *J Clin Invest* 1991; 87:1286-1294 and Alberts B, Bray D, Lewis J, Raff M, Roberts K, Watson JD, editors. Molecular biology of the cell. New York: Garland Publishing, Inc., 1989 with minor modifications. The liver homogenate is filtered
25 through silk to remove any remaining tissue debris and then is centrifuged at 10,000xg for 20 minutes at 40C. The supernatant is decanted and centrifuged at 100,000xg for 60 minutes at 40C. The pellet, microsomes and small vesicles, is resuspended and lightly homogenized in : 20 mM TRIS-HCl (pH 7.4), 50 mM 2-mercaptoethanol, 250 mM sucrose, 2 mM EDTA, 10 mM EGTA, 2 mM AEBSF, 0.1 mM TLCK, 0.1 mM
30 TPCK, 0.5 mM benzamidine, 25 ug/ml leupeptin, 5 ug/ml pepstatin A, 5 ug/ml;H5B antipain, 5 ug/ml chymostatin, 10 ug/ml aprotinin (Buffer A), to a final concentration of approximately 850 ug protein/ml. Protein concentration is determined by the Pierce Coomassie Plus Protein Assay using crystalline bovine serum albumin as a standard (Pierce Chemical Co., Rockford, IL).

35

- 21 -

Measurement of PTPase activity: The malachite green-ammonium molybdate method, as described by Lanzetta PA, Alvarez LJ, Reinach PS, Candia OA was used. An improved assay for nanomolar amounts of inorganic phosphate. *Anal. Biochem.* 1979;100:95-97, and adapted for the platereader, is used for the nanomolar detection of liberated phosphate by rat hepatic microsomal PTPases. The test procedure uses, as substrate, a dodecaphosphopeptide custom synthesized by AnaSpec, Inc. (San Jose, CA). The peptide, TRDIYETDYRK, corresponding to the 1142-1153 catalytic domain of the insulin receptor, is tyrosine phosphorylated on the 1146, 1150 and 1151 tyrosine residues. The microsomal fraction (83.25 ul) is preincubated for 10 min at 37deg.C with or without test compound (6.25ul) and 305.5 ul of the 81.83 mM HEPES reaction buffer, pH 7.4. Peptide substrate, 10.5 ul at a final concentration of 50 uM, is equilibrated to 37deg.C in a LABLINE Multi-Blok heater equipped with a titerplate adapter. The preincubated microsomal preparation (39.5 ul) with or without drug is added to initiate the dephosphorylation reaction, which proceeds at 37deg.C for 30 min. The reaction is terminated by the addition of 200 ul of the malachite green-ammonium molybdate-Tween 20 stopping reagent (MG/AM/Tw). The stopping reagent consists of 3 parts 0.45% malachite green hydrochloride, 1 part 4.2% ammonium molybdate tetrahydrate in 4 N HCl and 0.5% Tween 20. Sample blanks are prepared by the addition of 200 ul MG/AM/Tw to substrate and followed by 39.5 ul of the preincubated membrane with or without drug. The color is allowed to develop at room temperature for 30 min and the sample absorbances are determined at 650 nm using a platereader (Molecular Devices). Samples and blanks are prepared in quadruplicates. Screening activity of 50 uM (final) drug is accessed for inhibition of microsomal PTPases.

25

Calculations: PTPase activities, based on a potassium phosphate standard curve, are expressed as nmoles of phosphate released/min/mg protein. Test compound PTPase inhibition is calculated as percent of control. A four parameter non-linear logistic regression of PTPase activities using SAS release 6.08, PROC NLIN, is used for determining IC50 values of test compounds. All compounds were administered at a concentration of 50 uM. The following results were obtained using representative compounds of this invention:

30

- 22 -

Example	% Change from Control
1	-53
2	IC ₅₀ = 28.4 uM
3	IC ₅₀ = 29.9 uM
4	-11
5	-70
6	-71
7	-73
8	-28
9	-75
10	-72
11	-61
12	-50
13	-50
14	-55
15	-59
16	-53
17	-63
18	IC ₅₀ = 24.2 uM
19	IC ₅₀ = 21.4 uM
20	IC ₅₀ = 41.6 uM
21	IC ₅₀ = 35.5 uM
22	-55
25	IC ₅₀ = 17.9 uM
26	IC ₅₀ = 20.0 uM
27	IC ₅₀ = 24.9 uM
28	IC ₅₀ = 20.6 uM
29	-19
36	IC ₅₀ = 17.9 uM
37	IC ₅₀ = 36.8 uM
38	IC ₅₀ = 19.3 uM
phenylarsine (Reference)	-57

Inhibition of Tri-Phosphorylated Insulin Receptor Dodecaphosphopeptide
Dephosphorylation by hPTP1B

5 This standard pharmacological test procedure assess the inhibition of recombinant rat protein tyrosine phosphatase, PTP1B, activity using, as substrate, the phosphotyrosyl dodecapeptide corresponding to the 1142-1153 insulin receptor kinase domain, phosphorylated on the 1146, 1150 and 1151 tyrosine residues. The procedure used and results obtained are briefly described below.

10 Human recombinant PTP1B was prepared as described by Goldstein (see Goldstein et al. *Mol. Cell. Biochem.* **109**, 107, **1992**). The enzyme preparation used was in microtubes containing 500-700 µg/ml protein in 33 mM Tris-HCl, 2 mM EDTA, 10% glycerol and 10 mM 2-mercaptoethanol.

Measurement of PTPase activity. The malachite green-ammonium molybdate method, as described (Lanzetta et al. *Anal. Biochem.* **100**, 95, **1979**) and adapted for a platereader, is used for the nanomolar detection of liberated phosphate by recombinant PTP1B. The test procedure uses, as substrate, a dodecaphosphopeptide custom synthesized by AnaSpec, Inc. (San Jose, CA). the peptide, TRDIYETDYRK, corresponding to the 1142-1153 catalytic domain of the insulin receptor, is tyrosine phosphorylated on the 1146, 1150, and 1151 tyrosine residues. The recombinant rPTP1B is diluted with buffer (pH 7.4, containing 33 mM Tris-HCl, 2 mM EDTA and 50 mM b-mercaptoethanol) to obtain an approximate activity of 1000-2000 nmoles/min/mg protein. The diluted enzyme (83.25 mL) is preincubated for 10 min at 37°C with or without test compound (6.25 mL) and 305.5 mL of the 81.83 mM HEPES reaction buffer, pH 7.4 peptide substrate, 10.5 ml at a final concentration of 50 mM, and is equilibrated to 37°C. in a LABLINE Multi-Blok heater equipped with a titerplate adapter. The preincubated recombinant enzyme preparation (39.5 ml) with or without drug is added to initiate the dephosphorylation reaction, which proceeds at 37°C for 30 min. The reaction is terminated by the addition of 200 mL of the malachite green-ammonium molybdate-Tween 20 stopping reagent (MG/AM/Tw). The stopping reagent consists of 3 parts 0.45% malachite green hydrochloride, 1 part 4.2% ammonium molybdate tetrahydrate in 4 N HCl and 0.5% Tween 20. Sample blanks are prepared by the addition of 200 mL MG/AM/Tw to substrate and followed by 39.5 ml of the preincubated recombinant enzyme with or without drug. The color is allowed to develop at room temperature for 30 min. and the sample absorbances are determined at

- 24 -

650 nm using a platereader (Molecular Devices). Sample and blanks are prepared in quadruplicates.

- 5 Calculations: PTPase activities, based on a potassium phosphate standard curve, are expressed as nmoles of phosphate released/min/mg protein. Inhibition of recombinant PTP1B by test compounds is calculated as percent of phosphatase control. A four parameter non-linear logistic regression of PTPase activities using SAS release 6.08, PROC NLIN, is used for determining IC₅₀ values of test compounds. The following results were obtained.

10

Example	IC ₅₀ (μM)
1	0.322
2	0.025
3	0.029
4	-51 (2.5 uM)
5	-47 (2.5 uM)
6	-15 (2.5 uM)
7	-58 (2.5 uM)
8	0.74
9	2.19
10	0.44
11	0.27
12	0.176
13	0.095
14	0.68
15	0.348
16	0.397
17	1.32
18	0.401
19	0.123
20	0.077
21	0.118
22	0.118
23	-36 (2.5 uM)

- 25 -

Example	IC50 (μ M)
24	0.293
25	0.213
26	0.085
27	0.077
28	0.122
29	0.97
30	1.16
31	0.59
32	1.51
33	1.55
34	1.07
35	0.454
36	0.058
37	0.025
38	0.053
39	0.105
40	0.362
41	0.29
42	0.178
43	0.044
44	0.054
45	0.181
46	0.052
47	0.225
48	0.341
49	0.038
50	0.056
51	0.054
52	0.052
53	0.133
54	0.023
55	0.043

- 26 -

Example	IC50 (μ M)
56	0.047
57	0.028
58	0.025
59	0.08
60	0.071
61	0.100
62	0.105
63	0.369
64	0.512
65	0.074
66	0.788
67	0.103
68	0.231
69	-20 (1 μ M)
70	-41 (2.5 μ M)
71	-43 (2.5 μ M)
Phenylarsine oxide (reference standard)	39.7
Sodium orthovanadate (reference standard)	244.8
Ammonium molybdate tetrahydrate (reference standard)	8.7

The blood glucose lowering activity of representative compounds of this invention were demonstrated in an *in vivo* standard procedure using diabetic (ob/ob) mice. The procedures used and results obtained are briefly described below.

5 The non-insulin dependent diabetic (NIDDM) syndrome can be typically characterizes by obesity, hyperglycemia, abnormal insulin secretion, hyperinsulinemia and insulin resistance. The genetically obese-hyperglycemic ob/ob mouse exhibits many of these metabolic abnormalities and is thought to be a useful model to search for hypoglycemic agents to treat NIDDM [Coleman, D.: Diabetologia 14: 141-148, 1978].

10 In each test procedure, mice [Male or female ob/ob (C57 Bl/6J) and their lean littermates (ob/+ or +/+, Jackson Laboratories) ages 2 to 5 months (10 to 65 g)] of a similar age were randomized according to body weight into 4 groups of 10 mice. The mice were housed 5 per cage and are maintained on normal rodent chow with water ad

- 27 -

libitum. Mice received test compound daily by gavage (suspended in 0.5 ml of 0.5% methyl cellulose); dissolved in the drinking water; or admixed in the diet. The dose of compounds given ranges from 2.5 to 200 mg/kg body weight/day. The dose is calculated based on the fed weekly body weight and is expressed as active moiety. The positive control, ciglitazone (5-(4-(1-methylcyclohexylmethoxy)benzyl)-2,4-dione, see Chang, A., Wyse, B., Gilchrist, B., Peterson, T. and Diani, A. Diabetes 32: 830-838, 1983.) was given at a dose of 100 mg/kg/day, which produces a significant lowering in plasma glucose. Control mice received vehicle only.

On the morning of Day 4, 7 or 14 two drops of blood (approximately 50 ul) were collected into sodium fluoride containing tubes either from the tail vein or after decapitation. For those studies in which the compound was administered daily by gavage the blood samples were collected two hours after compound administration. The plasma was isolated by centrifugation and the concentration of glucose is measured enzymatically on an Abbott V.P. Analyzer.

For each mouse, the percentage change in plasma glucose on Day 4, 7 or 14 is calculated relative to the mean plasma glucose of the vehicle treated mice. Analysis of variance followed by Dunett's Comparison Test (one-tailed) are used to estimate the significant difference between the plasma glucose values from the control group and the individual compound treated groups (CMS SAS Release 5.18).

The results shown in the table below shows that the compounds of this invention are antihyperglycemic agents as they lower blood glucose levels in diabetic mice.

Example	Dose (mg/Kg)	% Change Glucose from Vehicle	% Change Insulin from Vehicle
1	100	-40.3	17a
2	25	-25.5	11.2a
3	25	-29.9	23.5a
10	100	-34.2	-51.9
11	100	-40.4	-44.9
12	100	-27.5	-15.8a
13	100	-36.7	5.0a
14	100	-39	19.8a
15	100	-35	4.1a
17	100	13.7a	-94.4

- 28 -

Example	Dose (mg/Kg)	% Change Glucose from Vehicle	% Change Insulin from Vehicle
19	100	-27.7	1.1a
24	25	-30.5	not determined
28	25	-20	not determined
42	25	-10.6	9.9a
48	25	4.7a	-30
55	25	-13.6	35.4a
Ciglitazone (reference standard)	100	-43	-39

a - no significant activity ($p < 0.05$) at this dose.

Based on the results obtained in the standard pharmacological test procedures, representative compounds of this invention have been shown to inhibit PTPase activity and lower blood glucose levels in diabetic mice, and are therefore useful in treating metabolic disorders related to insulin resistance or hyperglycemia, typically associated with obesity or glucose intolerance. More particularly, the compounds of this invention are useful in the treatment or inhibition of type II diabetes, and in modulating glucose levels in disorders such as type I diabetes. As used herein, the term modulating means maintaining glucose levels within clinically normal ranges.

Effective administration of these compounds may be given at a daily dosage of from about 1 mg/kg to about 250 mg/kg, and may be given in a single dose or in two or more divided doses. Such doses may be administered in any manner useful in directing the active compounds herein to the recipient's bloodstream, including orally, via implants, parenterally (including intravenous, intraperitoneal and subcutaneous injections), rectally, vaginally, and transdermally. For the purposes of this disclosure, transdermal administrations are understood to include all administrations across the surface of the body and the inner linings of bodily passages including epithelial and mucosal tissues. Such administrations may be carried out using the present compounds, or pharmaceutically acceptable salts thereof, in lotions, creams, foams, patches, suspensions, solutions, and suppositories (rectal and vaginal).

- 29 -

Oral formulations containing the active compounds of this invention may comprise any conventionally used oral forms, including tablets, capsules, buccal forms, troches, lozenges and oral liquids, suspensions or solutions. Capsules may contain mixtures of the active compound(s) with inert fillers and/or diluents such as the pharmaceutically acceptable starches (e.g. corn, potato or tapioca starch), sugars, artificial sweetening agents, powdered celluloses, such as crystalline and microcrystalline celluloses, flours, gelatins, gums, etc. Useful tablet formulations may be made by conventional compression, wet granulation or dry granulation methods and utilize pharmaceutically acceptable diluents, binding agents, lubricants, disintegrants, suspending or stabilizing agents, including, but not limited to, magnesium stearate, stearic acid, talc, sodium lauryl sulfate, microcrystalline cellulose, carboxymethylcellulose calcium, polyvinylpyrrolidone, gelatin, alginic acid, acacia gum, xanthan gum, sodium citrate, complex silicates, calcium carbonate, glycine, dextrin, sucrose, sorbitol, dicalcium phosphate, calcium sulfate, lactose, kaolin, mannitol, sodium chloride, talc, dry starches and powdered sugar. Oral formulations herein may utilize standard delay or time release formulations to alter the absorption of the active compound(s). Suppository formulations may be made from traditional materials, including cocoa butter, with or without the addition of waxes to alter the suppository's melting point, and glycerin. Water soluble suppository bases, such as polyethylene glycols of various molecular weights, may also be used.

It is understood that the dosage, regimen and mode of administration of these compounds will vary according to the malady and the individual being treated and will be subject to the judgment of the medical practitioner involved. It is preferred that the administration of one or more of the compounds herein begin at a low dose and be increased until the desired effects are achieved.

The following procedures describe the preparation of representative examples of this invention.

Example 1

(2S)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

Step a) *o*-Phenoxy-4-bromoacetophenone

Potassium carbonate (24.8 g, 179.8 mmol) was added into a mixture of 4-bromophenylacetyl bromide (50.0 g, 179.8 mmol), phenol (16.9 g, 179.8 mmol) and dry acetone (200mL). The reaction mixture was refluxed for 12 hours, cooled to room

- 30 -

temperature, poured into water, and extracted with ethyl ether. The organic extracts were dried over MgSO_4 . Evaporation gave a yellow solid (49.6 g, 94% yield): MS m/e 291 (M^+).

5 Step b) 3-(4-Bromophenyl)-1-benzofuran

A mixture of ω -phenoxy-4-bromoacetophenone (49.0 g, 167.8 mmol), polyphosphoric acid (100 g) and xylenes (300 mL) was refluxed for 12 hours. The reaction mixture cooled to room temperature, poured into water, and extracted with ethyl ether. The organic extracts were dried over MgSO_4 . Evaporation and purification
10 by flash chromatography on silica gel (hexanes/EtAOc 40:1) gave a yellow solid (36.2 g, 79% yield): mp 70-71 °C; MS m/e 272 (M^+);
Analysis for: $\text{C}_{14}\text{H}_9\text{BrO}$ Calc'd: C, 61.57; H, 3.32; Found: C, 61.80; H, 3.31

Step c) 3-(4'-Methoxy-biphenyl-4-yl)-benzofuran

15 4-Methoxy-benzeneboronic acid (14.17 g, 70.5 mmol) in ethyl alcohol (10 mL) was added into a mixture of 3-(4-bromophenyl)-1-benzofuran (17.5 g, 64.1 mmol), sodium carbonate (2N, 64.1 mL), tetrakis(triphenylphosphine)palladium(0) (2.23 g, 1.92 mmol), and toluene (200 mL). The reaction mixture was refluxed for 12 hours, cooled to room temperature, and treated with hydrogen peroxide (30%, 5 mL) for 1
20 hour. Then, the mixture was poured into water and extracted with ethyl acetate. The organic extracts were dried over MgSO_4 . Evaporation and crystallization from acetone/ethyl ether gave a white solid (14.9 g, 77% yield): mp 137-138 °C; MS m/e 300 (M^+);
Analysis for: $\text{C}_{21}\text{H}_{16}\text{O}_2$ Calc'd: C, 83.98; H, 5.37 Found: C, 83.70; H, 5.22

25

Step d) 4'-Benzofuran-3-yl-biphenyl-4-ol

Boron tribromide (1.0 M, 6.67 mL, 6.67 mmol) was added dropwise into a cold (-78 °C) mixture of 3-(4'-methoxy-biphenyl-4-yl)-benzofuran (2.0 g, 6.67 mmol), and dichloromethane (25 mL). The reaction mixture was allowed to come gradually to
30 room temperature and stirred for 10 hours. Then the mixture cooled to 0 °C and methyl alcohol (5 mL) was added dropwise. After stirring for 10 minutes the mixture was poured into water and extracted with ethyl ether. The organic extracts were dried over MgSO_4 . Evaporation and crystallization from ethyl ether/hexanes gave a yellow solid (1.69 g, 87% yield): mp 174-175; MS m/e 286 (M^+);
35 Analysis for: $\text{C}_{20}\text{H}_{14}\text{O}_2$ Calc'd: C, 83.90; H, 4.93 Found: C, 83.69; H, 4.88

- 31 -

Step e) [3-(4'-Hydroxy-biphenyl-4-yl)-benzofuran-2-yl]-phenyl-methanone

n-Butyllithium (2.5 N, 8.4 mL, 20.98 mmol) was added dropwise into a cold (-78 °C) mixture of 4'-benzofuran-3-yl-biphenyl-4-ol (3.0 g, 10.49 mmol) and tetrahydrofuran (50 mL). The mixture was allowed to gradually warm up to -40 °C and stirred for 3 hours. N-Methoxy N-methyl benzamide (1.6 mL, 10.49 mmol) was added dropwise into the mixture. The reaction mixture was allowed to gradually warm up to 0 °C and stirred for 30 minutes. The reaction was quenched with aqueous ammonium chloride, poured into water, acidified with HCl (2N), and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation and crystallization from ethyl ether/hexanes gave a yellow solid (2.9 g, 71% yield): mp 231-232; MS *m/e* 390 (M⁺);

Analysis for: C₂₇H₁₈O₃ Calc'd: C, 83.06; H, 4.65 Found: C, 82.63; H, 4.27

Step f) 4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-ol

Hydrazine monohydrate (1.38 g, 27.68 mmol) was added into a mixture of [3-(4'-hydroxy-biphenyl-4-yl)-benzofuran-2-yl]-phenyl-methanone (2.7 g, 6.92 mmol) and diethylene glycol (20 mL). The reaction mixture was stirred at 180 °C for 1 hour. The mixture cooled to room temperature and potassium hydroxide (1.16 g, 20.76 mmol) was gradually added. The mixture stirred at 130 °C for 10 hours, cooled to room temperature, poured into water, and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and crystallization from ethyl ether/hexanes/acetone gave a white solid (2.45 g, 94% yield): mp 151-153; MS *m/e* 376 (M⁺);

Analysis for: C₂₇H₂₀O₂ Calc'd: C, 86.15; H, 5.36 Found: C, 85.88; H, 5.13

Step g) (2S)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid methyl ester

Diethylazodicarboxylate (1.67 mL, 10.64 mmol) in benzene (20 mL) was added dropwise into a cold (0 °C) mixture of 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol (2.0 g, 5.32 mmol), (R)-(+)-3-phenyllactic acid methyl ester (1.91 g, 10.64 mmol), triphenylphosphine (2.8 g, 10.64 mmol) and benzene (50 mL). The reaction mixture was stirred at room temperature for 30 minutes, poured into water, and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and purification

- 32 -

by flash chromatography on silica gel (hexanes/EtAOc 8:1) gave a yellow oil (2.56 g, 89% yield): MS *m/e* 538 (M^+);

Analysis for: $C_{37}H_{30}O_4$ Calc'd: C, 82.51; H, 5.61 Found: C, 81.74; H, 5.78

5 Step h) (2S)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

Sodium hydroxide (2.5 N, 10 mL) was added into a mixture of (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid methyl ester (2.5 g, 4.65 mmol), methyl alcohol (40 mL) and tetrahydrofuran (40 mL). The reaction mixture was stirred for 1 hour, poured into water, acidified with HCl (2 N), and extracted with ethyl ether. The organic extracts were dried over $MgSO_4$. Evaporation and crystallization from ethyl ether/hexanes gave a white solid (2.32 g, 95% yield): mp 167-169; MS *m/e* 376 (M^+);

Analysis for: $C_{36}H_{28}O_4$ Calc'd: C, 82.42; H, 5.38 Found: C, 82.26; H, 5.36

15

Example 2

[(4,4''-Dimethoxy-5'-{2-(phenylmethyl)benzo[b]thien-3-yl}phenyl)[1,1':3',1''-terphenyl-2'-yl]oxy]-acetic acid

Step a) 3-(4'-Methoxy-biphenyl-4-yl)-benzo[b]thiophene

20 Palladium (II) acetate was added into a mixture of 3-bromo-benzo[b]-thiophene (1.4 g, 6.58 mmol), 4'-methoxy-biphenyl-4-boronic acid (1.5 g, 6.58 mmol), potassium carbonate (2.27 g, 16.45 mmol), acetone (20 mL) and H_2O (20 mL). The reaction mixture was stirred at 65 °C for 2h, poured into water, and extracted with ethyl acetate. The organic extracts were dried over $MgSO_4$. Evaporation and crystallization from ethyl ether/hexanes gave an off-white solid (1.76 g, 85% yield): mp 134-136; MS *m/e* 316 (M^+);

Analysis for: $C_{21}H_{16}OS$ Calc'd: C, 79.71; H, 5.10 Found: C, 78.98; H, 5.13

Step b) 4'-Benzo[b]thiophen-3-yl-biphenyl-4-ol

30 This compound was prepared from 3-(4'-methoxy-biphenyl-4-yl)-benzo[b]-thiophene, in substantially the same manner, as described in Example 1 step d, and was obtained as an off-white solid, mp 167-169 °C; MS *m/e* 302 (M^+);

Analysis for: $C_{20}H_{14}OS$ Calc'd: C, 79.44; H, 4.67 Found: C, 79.36; H, 4.52

- 33 -

Step c) [3-(4'-Hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-phenyl-methanone

This compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol, in substantially the same manner as described in Example 1 step e, and was obtained as a yellow solid, mp 205-207 °C; MS *m/e* 406 (M⁺);

5 Analysis for: C₂₇H₁₈O₂S Calc'd: C, 79.78; H, 4.46 Found: C, 78.95; H, 4.59

Step d) 4'-(2-Benzyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol

This compound was prepared from [3-(4'-hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-phenyl-methanone, in substantially the same manner as described in
10 Example 1 step f, and was obtained as a white solid, mp 178-180 °C; MS *m/e* 392 (M⁺);
Analysis for: C₂₇H₂₀OS Calc'd: C, 81.62; H, 5.14 Found: C, 81.60; H, 5.32

Step e) 4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-ol and
4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol

15 Bromine (1.47 mL, 28.69 mmol) in acetic acid (50 mL) was added dropwise over a 30 minutes period into a cold (5 °C) mixture of 4'-(2-benzyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol (7.5 g, 19.13 mmol), potassium acetate (18.6 g, 190.13 mmol), and acetic acid (200 mL). After the addition the mixture was poured into water, and extracted with ethyl ether. The organic extracts were washed with aqueous sodium
20 bisulfite and dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc/CH₂Cl₂ 3:1:1) gave 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-ol as a light yellow solid (2.4 g); mp 54-56 °C MS *m/e* 477 (M⁺);
Analysis for: C₂₇H₁₉BrOS Calc'd: C, 68.79; H, 4.06 Found: C, 68.37; H, 4.17

25 and 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol as a light yellow solid (4.7 g); mp 59-61 °C; MS *m/e* 548 (M⁺);
Analysis for: C₂₇H₁₈Br₂OS Calc'd: C, 58.93; H, 3.30 Found: C, 59.21; H, 3.57

Step f) 4,4''-Dimethoxy-5'-[4-[2-(phenylmethyl)benzo[b]thien-3-yl]phenyl]-
30 [1,1':3',1''-terphenyl]-2'-ol and 4-(2-benzyl-benzo[b]thiophen-3-yl)-5'-bromo-4''-
methoxy[1,1':3',1''-terphenyl]-4'-ol

Palladium (II) acetate (81 mg, 0.036 mmol) was added into a mixture of 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol (1.0 g, 1.82 mmol), 4-methoxy-benzeneboronic acid, barium hydroxide (0.93 g, 5.46 mmol), 1,2-
35 dimethoxyethane (10 mL), and water (10 mL). The mixture was stirred at 75 °C for 10 hours, poured into water, and extracted with ethyl ether. The organic extracts were

- 34 -

dried over MgSO_4 . Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 4:1) gave 4,4''-dimethoxy-5'-{4-[2-(phenylmethyl)benzo[b]thien-3-yl]phenyl}[1,1';3',1''-terphenyl]-2'-ol as a yellow solid (0.45 g.); mp 80-82; MS m/e 604 (M^+);

5 Analysis for: $\text{C}_{41}\text{H}_{32}\text{O}_3\text{S}$ Calc'd: C, 81.43; H, 5.33 Found: C, 80.19; H, 5.44

and 4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-4''-methoxy[1,1';3',1''-terphenyl]-4'-ol as a yellow solid (0.45 g.); mp 74-76; MS m/e 576 (M^+);

Analysis for: $\text{C}_{34}\text{H}_{25}\text{O}_2\text{S}$ Calc'd: C, 70.71; H, 4.36 Found: C, 69.78; H, 5.31

10

Step g) [(4,4''-Dimethoxy-5'-(2-(phenylmethyl)benzo[b]thien-3-yl)phenyl)-[1,1':3',1''-terphenyl]-2'-yl)oxy]-acetic acid

Methyl bromoacetate (0.16 mL, 1.66 mmol) was added dropwise into a mixture of 4,4''-dimethoxy-5'-{4-[2-(phenylmethyl)benzo[b]thien-3-yl]phenyl}[1,1';3',1''-terphenyl]-2'-ol (1.0 g, 1.66 mmol), potassium carbonate (0.23 g, 1.66 mmol), and N,N-dimethylformamide (10 mL). The reaction mixture was stirred at 75 °C for 2 hours, and then poured into water and extracted with ethyl acetate. The organic extracts were dried over MgSO_4 . Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 4:1) gave a yellow oil (1.05 g), which was taken in methyl alcohol (10 mL) and tetrahydrofuran (10 mL), and treated with NaOH (2.5 N 3.0 mL) for 30 minutes. The new reaction mixture was poured into water, acidified with HCl (2 N), and extracted with ethyl ether. The organic extracts were dried over MgSO_4 . Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 3:1) gave white solid (0.79 g, 72 % yield): mp 99-101; MS m/e 662 (M^+);

25 Analysis for: $\text{C}_{43}\text{H}_{34}\text{O}_5\text{S}$ Calc'd: C, 77.92; H, 5.17 Found: C, 77.03; H, 5.29

Example 3

[4-(2-Benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-4''-methoxy[1,1':3',1''-terphenyl]-4'-yl)oxy]-acetic acid

30 The title compound was prepared from 4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-4''-methoxy[1,1';3',1''-terphenyl]-4'-ol, in substantially the same manner, as described in Example 2 step g, and was obtained as a white solid, mp 94-96 °C; MS m/e 634 (M^+);

Analysis for: $\text{C}_{36}\text{H}_{27}\text{BrO}_4\text{S}$ Calc'd: C, 68.03; H, 4.28 Found: C, 67.57; H, 4.29

35

- 35 -

Example 4**(4'-Benzofuran-3-yl-biphenyl-4-yloxy)-acetic acid**

Sodium hydride (0.15 g, 3.49 mmol) was added portionwise into a mixture of 4'-benzofuran-3-yl-biphenyl-4-ol (1.0 g, 3.49 mmol), and N,N-dimethylformamide (10 mL). The reaction mixture was stirred for 1 hour, and then methyl bromoacetate (0.49 mL, 5.23 mmol) was added dropwise. The mixture was stirred for 30 minutes, poured into water, and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation gave a yellow oil (1.25 g). The residue was taken in methyl alcohol (20 mL) and tetrahydrofuran (20 mL), and treated with sodium hydroxide (2.5 N, 5.0 mL) for 30 minutes. The mixture was then poured into water, acidified with HCl (2 N) and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and crystallization from ethyl ether/hexanes gave a white solid (0.38 g, 32% yield): mp 196-197; MS *m/e* 344 (M⁺); Analysis for: C₂₂H₁₆O₄ Calc'd: C, 76.73; H, 4.68 Found: C, 76.63; H, 4.58

Example 5**2-(4'-Benzofuran-3-yl-biphenyl-4-yloxy)-3-phenyl-propionic acid**

The title compound was prepared from 4'-benzofuran-3-yl-biphenyl-4-ol, in substantially the same manner, as described in Example 2 step g, and was obtained as a white solid, mp 142-143 °C; MS *m/e* 434 (M⁺); Analysis for: C₂₉H₂₂O₄ Calc'd: C, 80.17; H, 5.10 Found: C, 80.54; H, 5.09

Example 6**[4'-(2-Bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid****25 Step a) 2-Bromo-3-(4'-methoxy-biphenyl-4-yl)-benzofuran**

Bromine (0.29 mL, 5.67 mmol) in acetic acid (50 mL) was added dropwise over a 30 minutes period into a cold (5 °C) mixture of 3-(4'-methoxy-biphenyl-4-yl)-benzofuran (1.7 g, 5.67 mmol), potassium acetate (5.55 g, 56.7 mmol), and acetic acid (20 mL). After the addition, the mixture was poured into water, and extracted with ethyl ether. The organic extracts were washed with aqueous sodium bisulfite and dried over MgSO₄. Evaporation and crystallization gave an off-white solid (1.69 g, 79% yield): mp 136-137 °C MS *m/e* 378 (M⁺); Analysis for: C₂₁H₁₅BrO₂ Calc'd: C, 66.51; H, 3.99 Found: C, 66.17; H, 3.84

- 36 -

Step b) 4'-(2-Bromo-benzofuran-3-yl)-biphenyl-4-ol

This compound was prepared from 2-bromo-3-(4'-methoxy-biphenyl-4-yl)-benzofuran, in substantially the same manner, as described in Example 1 step d, and was obtained as a white solid, mp 150-151 °C; MS *m/e* 364 (M^+);

5 Analysis for: $C_{20}H_{13}BrO_2$ Calc'd: C, 65.77; H, 3.59 Found: C, 65.47; H, 3.46

Step c) [4'-(2-Bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid

This compound was prepared from 4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-ol, in substantially the same manner, as described in Example 4, and was obtained as a

10 white solid, mp 178-180 °C; MS *m/e* 423 (M^+);

Analysis for: $C_{22}H_{15}BrO_4$ Calc'd: C, 62.43; H, 3.57 Found: C, 61.68; H, 3.42

Example 72-[4'-(2-Bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

15 The title compound was prepared from 4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-ol and 3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 148-149 °C; MS *m/e* 512 (M^+);

Analysis for: $C_{29}H_{21}BrO_4$ Calc'd: C, 67.85; H, 4.12 Found: C, 67.68; H, 4.31

20

Example 84'-(2-Butyl-benzofuran-3-yl)-biphenyl-4-ol

The title compound was prepared from 4'-benzofuran-3-yl-biphenyl-4-ol and N-methoxy-N-methyl-n-butylamide, in substantially the same manner, as described in

25 Example 1, steps e-f, and was obtained as a white solid, mp 128-129 °C; MS *m/e* 342 (M^+);

Analysis for: $C_{24}H_{22}O_2$ Calc'd: C, 84.18; H, 9.48 Found: C, 84.16; H, 6.45

Example 930 [4'-(2-Butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid

The title compound was prepared from 4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-ol, in substantially the same manner, as described in Example 4, and was obtained as a white solid, mp 126-128 °C; MS *m/e* 400 (M^+);

Analysis for: $C_{26}H_{24}O_4$ Calc'd: C, 77.98; H, 6.04 Found: C, 77.80; H, 6.06

35

- 37 -

Example 10**2-[4'-(2-Butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

The title compound was prepared from 4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-ol, and 3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 169-170 °C; MS *m/e* 491 (M+H)⁺;

Analysis for: C₃₃H₃₀O₄ Calc'd: C, 80.79; H, 6.16 Found: C, 80.75; H, 6.12

Example 11**2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, and 3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 164-166 °C; MS *m/e* 524 (M⁺);

Analysis for: C₃₆H₂₈O₄ Calc'd: C, 82.42; H, 5.38 Found: C, 82.14; H, 5.20

Example 12**2-[4'-(2-Butyl-benzob[thiopen-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

The title compound was prepared from 4'-(2-butyl-benzo[b]-thiophen-3-yl)-biphenyl-4-ol, and 3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 159-161 °C; MS *m/e* 507 (M+H)⁺;

Analysis for: C₃₃H₃₀O₃S Calc'd: C, 78.23; H, 5.97 Found: C, 78.12; H, 6.02

Example 13**2-[4'-(2-Benzyl-benzob[thiophen-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

The title compound was prepared from 4'-(2-benzyl-benzo[b]-thiophen-3-yl)-biphenyl-4-ol, and 3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 165-167 °C; MS *m/e* 540 (M⁺);

Analysis for: C₃₆H₂₈O₃S Calc'd: C, 79.97; H, 5.22 Found: C, 79.96; H, 5.38

Example 14**2-[4'-(2-Benzoyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

The title compound was prepared from 4'-(2-benzoyl-benzofuran-3-yl)-biphenyl-4-ol, and 3-phenyllactic acid methyl ester, in substantially the same manner,

- 38 -

as described in Example 1, steps g-h, and was obtained as a yellow solid, mp 159-161°C; MS *m/e* 538 (*M*⁺);

Analysis for: C₃₆H₂₆O₅ Calc'd: C, 80.28; H, 4.86 Found: C, 80.15; H, 4.91

5

Example 15

(2R)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 167-169 °C;

10 MS *m/e* 524 (*M*⁺);

Analysis for: C₃₆H₂₈O₄ Calc'd: C, 82.42; H, 5.38 Found: C, 82.43; H, 5.53

Example 16

(2S)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic tromethamine salt

15

A mixture of (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid (1.0 g, 1.91 mmol), tromethamine (0.23 g, 1.91 mmol), tetrahydrofuran (10 mL), and water (1.0 mL) was stirred at 60 °C for 1 hour. The volatiles were removed in vacuo and the residue was washed with water and dried to give a white solid (1.1g, 89% yield); mp 147-148; MS *m/e* 523 (*M-H*)⁺;

20

Analysis for: C₃₆H₂₇O₄.tromethamine x 1.5 H₂O Calc'd: C, 71.41; H, 6.29; N, 2.08 Found: C, 71.33; H, 6.35; N, 2.40

Example 17

25

(2R)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-propionic acid

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, and (S)-(+)- lactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 112-114 °C; MS *m/e* 448 (*M*⁺);

30

Analysis for: C₃₀H₂₄O₄ Calc'd: C, 80.34; H, 5.39 Found: C, 79.86; H, 5.73

Example 18

(2R)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid

35

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, and (S)-(+)- mandelic acid methyl ester, in substantially the same manner, as

- 39 -

described in Example 1, steps g-h , and was obtained as a white solid, mp 177-179 °C; MS *m/e* 509 (M-H)⁺;

Analysis for: C₃₅H₂₆O₄ x 0.4 H₂O Calc'd: C, 81.19; H, 5.22 Found: C, 81.34; H, 5.46

5

Example 19

(2R)-2-{4'-[2-(4-Fluoro-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

Step a) (4-Fluoro-phenyl)-[3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-methanol

10

This compound was prepared from 3-(4'-methoxy-biphenyl-4-yl)-benzo[b]-thiophene, and 4-fluoro-benzaldehyde, in substantially the same manner, as described in Example 1 , step e, and was obtained as a white solid, mp 66-68 °C; MS *m/e* 440 (M⁺);

Analysis for: C₂₈H₂₁FO₂S Calc'd: C, 76.34; H, 4.81 Found: C, 75.82; H, 4.89

15

Step b) 2-(4-Fluoro-benzyl)-3-(4'-methoxy-biphenyl-4-yl)-benzo[b]-thiophene

Trifluoroacetic acid (5 mL) was added dropwise into a mixture of (4-fluoro-phenyl)-[3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-methanol (1.2 g, 2.72 mmol), sodium borohydride (0.51 g, 13.75 mmol), and ethyl ether (10 mL). The reaction mixture was stirred for 30 minutes, poured into water, basified with sodium hydroxide (2 N), and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and crystallization from ethyl ether/hexanes gave a white solid (1.02 g, 88% yield): mp 100-102; MS *m/e* 424 (M⁺);

25

Step c) (2R)-2-{4'-[2-(4-Fluoro-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

This compound was prepared from 2-(4-fluoro-benzyl)-3-(4'-methoxy-biphenyl-4-yl)-benzo[b]-thiophene, in substantially the same manner, as described in Example 1 step d, and g-h , and was obtained as a white solid, mp 151-153 °C; MS *m/e* 558 (M⁺);

30

Analysis for: C₃₆H₂₆FO₃S Calc'd: C, 77.40; H, 4.87 Found: C, 77.55; H, 4.58

- 40 -

Example 20**(2R)-2-[4'-(2-(4-Methoxy-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy)-3-phenyl-propionic acid**

5 The title compound was prepared from 3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophene, and 4-methoxy-benzaldehyde in substantially the same manner, as described in Example 19, and was obtained as a white solid, mp 75-77 °C; MS *m/e* 569 (M-H)⁺;

Analysis for: C₃₇H₃₀O₄S Calc'd: C, 77.87; H, 5.30 Found: C, 76.57; H, 5.39

10

Example 21**[4'-(2-Butyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid**

The title compound was prepared from 4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-ol, and mandelic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 157-159 °C; MS *m/e* 493 (M+H)⁺;

15

Analysis for: C₃₂H₂₈O₃S Calc'd: C, 78.02; H, 5.73 Found: C, 77.67; H, 5.91

Example 22

20 **(2R)-2-[4'-(2-(Hydroxy-phenyl-methyl)-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid**

Sodium borohydride (0.15 g, 4.06 mmol) was added portionwise into a cold (0°C) mixture of (2R)-2-[4'-(2-benzoyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid methyl ester (1.5, 2.7 mmol), methyl alcohol (20 mL) and tetrahydrofuran (5 mL). The reaction mixture was allowed to come to room temperature and stirred for 30 minutes. The mixture was then poured into water, acidified with HCl (2 N), and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation gave a yellow oil (1.4 g), which was taken in methyl alcohol (20 mL), tetrahydrofuran (20 mL), and treated with sodium hydroxide (2.5 N, 5.0 mL) for 30 minutes. The mixture was poured into water, acidified with HCl (2 N) and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 5:1) gave an off-white solid (1.16 g, 79 % yield): mp 95-97; MS *m/e* 539 (M-H)⁺;

25

30

Analysis for: C₃₆H₂₈O₅ Calc'd: C, 79.98; H, 5.22 Found: C, 79.92; H, 5.34

- 41 -

Example 23

(2S)-2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid methyl ester

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, in substantially the same manner, as described in Example 1, step g, and was obtained as a white solid, mp 67-69 °C; MS *m/e* 538 (M⁺);

Analysis for: C₃₇H₃₀O₄ Calc'd: C, 82.51; H, 5.61 Found: C, 81.74; H, 5.78

Example 24

10 2-[4'-(2-Benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid
Step a) 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid ethyl ester

2-Bromo-propionic acid ethyl ester (1.28 mL, 9.84 mmol) was added dropwise into a mixture of 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol (3.7 g, 9.48 mmol), potassium carbonate (1.49 g, 10.82 mmol), and N,N-dimethylformamide (10 mL). The reaction mixture was stirred at 70 °C for 10 hours, poured in to water, acidified with HCl (2 N), and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 10:1) gave a viscous clear oil (4.2 g, 96 % yield): MS *m/e* 476 (M⁺);

15
20 Analysis for: C₃₂H₂₈O₄ x 0.2 H₂O Calc'd: C, 80.04; H, 5.96 Found: C, 80.03; H, 6.01

Step b) 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid

Lithium bis(trimethylsilyl)amide (1.0 M, 1.68 mL, 1.68 mmol) was added dropwise into a cold (-78 °C) mixture of 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid ethyl ester (0.8 g, 1.68 mmol), and tetrahydrofuran (8.0 mL). The mixture was allowed to gradually warm up to -45 °C, and stirred for 2 hours. Benzyl bromide (0.29 mL, 2.52 mmol) was added into the reaction mixture. The temperature was gradually raised to room temperature and the reaction was stirred for 15 hours. The mixture was then poured in to water, acidified with HCl (2 N) to pH = 5, and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 5:1) gave an clear oil (0.43 g). The product was saponified according to Example 1 step h, to give a white solid (0.32 g, 34% yield for two steps): MS *m/e* 538 (M⁺);

25
30
35 Analysis for: C₃₇H₃₀O₄ Calc'd: C, 81.68; H, 5.67 Found: C, 81.66; H, 5.69

Example 25

(2R)-2-{4'-[2-(3,4-Dimethoxy-benzyl)-benzo[b]thiophen-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid

5 The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol, and 3,4-dimethoxy-benzaldehyde in substantially the same manner, as described in Example 20, and was obtained as a white solid, mp 59-61 °C; MS *m/e* 599 (M-H)⁺;

10 Analysis for: C₃₈H₂₂O₅S x 1.5 H₂O Calc'd: C, 72.70; H, 5.62 Found: C, 72.40; H, 5.50

Example 26

(2R)-2-{4'-[2-(2,4-Dimethoxy-benzyl)-benzo[b]thiophen-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid

15 The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol, and 2,4-dimethoxy-benzaldehyde in substantially the same manner, as described in Example 20, and was obtained as a white solid, mp 70-72 °C; MS *m/e* 599 (M-H)⁺;

20 Analysis for: C₃₈H₂₂O₅S x 0.5 H₂O Calc'd: C, 74.85; H, 5.37 Found: C, 74.80; H, 5.44

Example 27

(2R) 2-{4'-[2-(2,2-Dimethyl-benzo[1,3]dioxol-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid

25

Step a) 4-[3-(4'-Hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-ylmethyl]-benzene-1,2-diol

30 This compound was prepared from 4'-[2-(2,4-dimethoxy-benzyl)-benzo[b]thiophen-3-yl]-biphenyl-4-ol, in substantially the same manner, as described in Example 1 step d, and was obtained as a white solid, mp 148-150 °C; MS *m/e* 452 (M⁺);

Analysis for: C₂₉H₂₄O₃S x 0.5 H₂O Calc'd: C, 75.46; H, 5.46 Found: C, 75.53; H, 5.41

- 43 -

Step b) 4'-[2-(2,2-Dimethyl-benzo[1,3]dioxol-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-ol

Camphorsulfonic acid (0.2 g) was added into a mixture of 4-[3-(4'-hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-ylmethyl]-benzene-1,2-diol (1.4 g, 3.3 mmol), and 2,2-dimethoxy propane (5.0 mL). The reaction mixture was stirred at 40 °C for 24 hours, poured into water, and extracted with ethyl ether. The organic extracts were washed with aqueous sodium bicarbonate, and then dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 4:1) gave a gray solid (1.19 g, 72% yield): mp 58-60 °C; MS *m/e* 464 (M⁺);

10 Analysis for: C₃₀H₂₄O₃S x 1 H₂O Calc'd: C, 74.66; H, 5.43 Found: C, 74.53; H, 5.74

Step c) (2R) 2-{4'-[2-(2,2-Dimethyl-benzo[1,3]dioxol-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid

This compound was prepared from 4'-[2-(2,2-dimethyl-benzo[1,3]dioxo-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester in substantially the same manner, as described in Example 1 steps g-h , and was obtained as a white solid, mp 68-70 °C; MS *m/e* 611 (M-H)⁺;

15 Analysis for: C₃₉H₃₂O₅S x 0.5 H₂O Calc'd: C, 75.34; H, 5.35 Found: C, 75.35; H, 5.35

20

Example 28

(2R)-2-{4'-[2-(3,4-Dihydroxy-benzyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid

A mixture of (2R) 2-{4'-[2-(2,2-dimethyl-benzo[1,3]dioxol-5-ylmethyl)-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy}-3-phenyl-propionic acid (0.38 g, 0.62 mmol), tetrahydrofuran (30 mL) and HCl (2 N, 5.0 mL) was stirred at 50 °C for 10 hours. The reaction mixture was then poured into water, and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on acidic silica gel (hexanes/EtAOc 2:1) gave a pink solid (0.26 g, 72% yield): mp 108-110 °C; MS *m/e* 571 (M-H)⁺;

25

30 Analysis for: C₃₆H₂₈O₅S x 0.5 H₂O Calc'd: C, 74.33; H, 5.03 Found: C, 74.22; H, 5.28

Example 29(2R)-2-[4'-(2-Benzyl-thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acidStep a) 3-(4'-Methoxy-biphenyl-4-yl)-thiophene-2-carboxaldehyde

This compound was prepared from 3-bromo-thiophene-2-carboxaldehyde and 4'-methoxy-biphenyl-4-boronic acid in substantially the same manner, as described in Example 2 step a, and was obtained as an off-white solid, mp 139-141 °C; MS *m/e* 294 (M⁺);

Analysis for: C₁₈H₁₄O₂S Calc'd: C, 73.44; H, 4.79 Found: C, 73.37; H, 4.77

10 Step b) [3-(4'-Methoxy-biphenyl-4-yl)-thiophen-2-yl]-phenyl-methanol

Phenyl magnesium bromide (2 M, 13.6 mL, 27.21 mmol) was added dropwise into a cold (0 °C) mixture of 3-(4'-methoxy-biphenyl-4-yl)-thiophene-2-carboxaldehyde (8.0 g, 27.21 mmol), and tetrahydrofuran (100 mL). The reaction mixture was stirred for 30 minutes, and then quenched with aqueous ammonium chloride, poured into water and extracted with ethyl acetate. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 4:1) gave an off-white solid (7.2 g, 71% yield): mp 115-117 °C; MS *m/e* 372 (M⁺);

Analysis for: C₂₄H₂₀O₂S Calc'd: C, 77.39; H, 5.41 Found: C, 77.12; H, 5.48

20 Step c) (2R)-2-[4'-(2-Benzyl-thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

This compound was prepared from [3-(4'-methoxy-biphenyl-4-yl)-thiophen-2-yl]-phenyl-methanol in substantially the same manner, as described in Example 19 step b, and Example 1 steps d-h, and was obtained as a white solid, mp 152-154 °C; MS *m/e* 489 (M-H)⁺;

Analysis for: C₃₂H₂₆O₃S Calc'd: C, 78.34; H, 5.34 Found: C, 78.32; H, 5.52

Example 3030 3-Phenyl-2-[4'-(2-thiazole-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acidStep a) 4'-(2-Thiazole-2-yl-methyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol

This compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol and thiazole-2-carboxaldehyde in substantially the same manner, as described in Example 1 step e, and Example 19 step b (with the modification of using triethylsilane in place of sodium borohydride as the reducing agent), and was obtained as a light yellow solid, mp 208-210 °C; MS *m/e* 399 (M⁺);

- 45 -

Analysis for: $C_{24}H_{17}NOS_2 \times 0.5 H_2O$ Calc'd: C, 70.56; H, 4.44; N, 3.43 Found: C, 70.60; H, 4.36; N, 3.35

5 Step b) 3-Phenyl-2-[4'-(2-thiazole-2-yl)methyl-benzo[b]thiophene-3-yl]-biphenyl-4-yloxy]-propionic acid

This compound was prepared from 4'-(2-thiazole-2-yl-methyl-benzo[b]-thiophen-3-yl)-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 85-88 °C; MS *m/e* 547 (M^+);

10 Analysis for: $C_{33}H_{25}NO_3S_2 \times 1 H_2O$ Calc'd: C, 70.06; H, 4.81; N, 2.48 Found: C, 69.81; H, 4.80; N, 2.68

Example 31

15 (2S)-2-[4'-(2-Benzyl-furo[2,3]pyridin-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

Step a) 4'-(2-Benzyl-furo[2,3-b]pyridin-3-yl)-biphenyl-4-ol

This compound was prepared from 3-bromo-2-benzyl-furo[2,3-b]pyridine, in substantially the same manner, as described in Example 2, steps e and a, and Example 1 step d, and was obtained as an off-white solid, mp 185-186 °C; MS *m/e* 386 (M^+);

20 Analysis for: $C_{26}H_{19}NO_2 \times 0.4 H_2O$ Calc'd: C, 81.18; H, 5.19; N, 3.64 Found: C, 81.18; H, 5.14; N, 3.47

Step b) (2S)-2-[4'-(2-Benzyl-furo[2,3]pyridin-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid potassium salt

25 This compound was prepared from 4'-(2-benzyl-furo[2,3-b]pyridin-3-yl)-biphenyl-4-ol, and (R)-(+)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h. In step h the hydrolysis was done with potassium hydroxide, and removal of the solvents gave the desired product. The product was obtained as an off-white solid, mp 85-88 °C; MS *m/e* 593 (M^+);

30 Analysis for: $C_{33}H_{26}NO_4K \times 1.2 H_2O$ Calc'd: C, 71.82; H, 4.89; N, 2.39 Found: C, 71.84; H, 4.93; N, 2.13

- 46 -

Example 32

(2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid sodium salt

5 The title compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol and pyridine-2 carboxaldehyde, in substantially the same manner, as described in Example 2, steps e and a, and Example 30, and was obtained as a white solid; MS *m/e* 540 (M-H)⁺;

Analysis for: C₃₃H₂₆NO₃SNa x 1 H₂O Calc'd: C, 72.29; H, 4.81; N, 2.41 Found: C, 72.67; H, 4.66; N, 2.41

10

Example 33

(2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid

15 The title compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol and pyridine-2 carboxaldehyde, in substantially the same manner, as described in Example 2, steps e and a, and Example 30, and was obtained as a white solid; MS *m/e* 540 (M-H)⁺;

Analysis for: C₃₃H₂₇NO₃S x 0.5 H₂O Calc'd: C, 76.36; H, 5.09; N, 2.54 Found: C, 776.46; H, 5.14; N, 2.29

20

Example 34

4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-ol

25 The title compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol in substantially the same manner, as described in Example 1 steps e, was and obtained as a light yellow solid (2.4 g); mp 54-56 °C MS *m/e* 477 (M⁺);

Analysis for: C₂₇H₁₉BrOS Calc'd: C, 68.79; H, 4.06 Found: C, 68.37; H, 4.17

Example 35

4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol

30 The title compound was prepared from 4'-benzo[b]thiophen-3-yl-biphenyl-4-ol in substantially the same manner, as described in Example 1 steps e, was and obtained as a light yellow solid (4.7 g); mp 59-61 °C; MS *m/e* 548 (M⁺);

Analysis for: C₂₇H₁₈Br₂OS Calc'd: C, 58.93; H, 3.30 Found: C, 59.21; H, 3.57

- 47 -

Example 36

(2R)-2-[4'-(2-Benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid

5 The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3-bromo-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 81-83 °C; MS *m/e* 618 (M⁺);

Analysis for: C₃₆H₂₇BrO₃S Calc'd: C, 69.79; H, 4.39 Found: C, 69.39; H, 4.40

10

Example 37

(2R)-2-[4'-(2-Benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid

15 The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 87-89 °C; MS *m/e* 696 (M⁺);

Analysis for: C₃₆H₂₆Br₂O₃S Calc'd: C, 61.91; H, 3.75 Found: C, 69.99; H, 3.86

Example 38

20 (2R)-2-[4-(2-Benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid

The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was
25 obtained as a white solid, mp 82-84 °C; MS *m/e* 645 (M-H)⁺;

Analysis for: C₄₃H₃₄O₄S Calc'd: C, 79.85; H, 5.30 Found: C, 79.67; H, 5.79

Example 39

[4'-(2-Benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-acetic acid

30 The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2 step g, and was obtained as an off-white solid, mp 183-185 °C; MS *m/e* 606 (M⁺);

Analysis for: C₂₉H₂₀Br₂O₃S Calc'd: C, 57.26; H, 3.31 Found: C, 57.42; H, 3.53

35

- 48 -

Example 40[4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-yloxy]-acetic acid

The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2 step g, and was obtained as a white solid, mp 150-152 °C; MS *m/e* 528 (M⁺);

Analysis for: C₂₉H₂₀Br₂O₃S Calc'd: C, 65.79; H, 4.00 Found: C, 65.50; H, 4.29

Example 41

10 (2S)-2-[4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-phenyl-butyric acid

The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol, and (R)-(-)-2-hydroxy-4-phenylbutyric acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 85-87 °C; MS *m/e* 709 (M-H)⁺;

Analysis for: C₃₇H₂₈Br₂O₃S Calc'd: C, 62.37; H, 3.96 Found: C, 62.03; H, 4.06

Example 424-[4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-yloxy]-butyric acid

20 The title compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3-bromo-biphenyl-4-ol, and 4-bromobutyric acid methyl ester, in substantially the same manner, as described in Example 2 step g, and was obtained as a white solid, mp 135-137 °C; MS *m/e* 555 (M-H)⁺;

Analysis for: C₃₁H₂₅BrO₃S x 0.3 H₂O Calc'd: C, 66.01; H, 4.54 Found: C, 66.13; H, 4.78

Example 43(2R)-2-[4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid

30 Step a) (2R)-2-[4'-(2-Benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid methyl ester

This compound was prepared from 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-ol, and (S)-(+)-a-1,3-dioxo-2-isoindolinebutyric acid methyl ester, in substantially the same manner, as described in Example 1 step g, and was obtained as a light yellow oil (1.2 g, 83% yield); MS *m/e* 793 (M⁺);

- 49 -

Step b) (2R)-2-[4'-(2-Benzyl-benzo[b]-thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid

Trimethylsilyl iodide (0.29 mL, 2.07 mmol) was added into a mixture of (2R)-2-[4'-(2-benzyl-benzo[b]-thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid methyl ester (1.1 g, 1.38 mmol), and methylene chloride (10 mL). The reaction mixture was stirred at 50 °C for 24 hours. An additional 0.29 mL of trimethylsilyl iodide was added and the mixture was stirred for 48 hours, poured into water and extracted with ethyl acetate. The organic extracts were washed with aqueous sodium bisulfite and dried over MgSO₄. Evaporation and purification by flash chromatography on silica gel (hexanes/EtAOc 2:1) gave white solid (0.92 g, 94 % yield): mp 113-115 °C; MS *m/e* 778 (M-H)⁺;

Analysis for: C₃₉H₂₇Br₂NO₅S Calc'd: C, 59.94; H, 3.48; N, 1.79 Found: C, 59.98; H, 3.77; N, 1.74

15

Example 44

N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-methoxycarbonyl-propyl]-phthalamic acid

Potassium carbonate (0.01 g) was added into a mixture of (2R)-2-[4'-(2-benzyl-benzo[b]-thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid methyl ester (0.5, 0.063 mmol), methyl alcohol (10 mL), and water (0.5 mL). The reaction mixture was stirred at room temperature for 24 hours, poured into water, acidified with HCl (2 N), and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on acidic silica gel (hexanes/EtAOc 4:1) gave white solid (0.32 g, 63% yield):

mp 95-97 °C; MS *m/e* 811 (M⁺);

Analysis for: C₄₀H₃₁Br₂NO₆S Calc'd: C, 59.04; H, 3.81; N, 1.72 Found: C, 58.43; H, 4.09; N, 1.59

30

Example 45

N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-carboxy-propyl]-phthalamic acid

Sodium hydroxide (2 N, 5.0 mL) was added into a mixture of (2R)-2-[4'-(2-benzyl-benzo[b]-thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid methyl ester (1.0g, 1.26 mmol), methyl alcohol (10 ml) and tetrahydrofuran (10 mL). The reaction mixture was stirred at room temperature

- 50 -

for 1 hour, poured into water, acidified with HCl (2 N), and extracted with ethyl ether. The organic extracts were dried over MgSO₄. Evaporation and purification by flash chromatography on acidic silica gel (hexanes/EtAc 2:1) gave white solid (0.82 g, 74% yield): mp 130-132 °C; MS *m/e* 797 (M⁺);

- 5 Analysis for: C₃₉H₂₉Br₂NO₆S Calc'd: C, 58.59; H, 3.66 ;N, 1.75 Found: C, 58.20; H, 3.97; N, 1.67

Example 46

(2R)-2-[4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-chloro-[1,1':3'.1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid

- 10 This compound was prepared from 4-(2-benzyl-benzo[b]-thiophen-3-yl)-4''-chloro[1,1':3'.1'']-terphenyl-4'-ol , and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h , and was obtained as a white solid, mp 85-87 °C; MS *m/e* 649 (M-H)⁺;
Analysis for: C₄₂H₃₁ClO₃S x 1.2 H₂O Calc'd: C, 74.97; H, 5.00 Found: C, 74.74; H,
15 4.84

Example 47

(2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-phenyl-butyric acid

- 20 The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol, and (R)-(-)-2-hydroxy-4-phenylbutyric acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h , and was obtained as a white solid, mp 152-154 °C; MS *m/e* 537 (M-H)⁺;
Analysis for: C₃₇H₃₀O₄ Calc'd: C, 82.50; H, 5.61 Found: C, 82.26; H, 5.56

25

Example 48

(2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid

- 30 The title compound was prepared from 4'-(2-benzyl-benzofuran -3-yl)-biphenyl-4-ol, and (S)-(+)-a-1,3-dioxo-2-isoindolinebutyric acid methyl ester, in substantially the same manner, as described in Example 1 step g, and Example 43 step b, and was obtained as a yellow solid, mp 182-184 °C; MS *m/e* 606 (M-H)⁺;
Analysis for: C₃₉H₂₉NO₆ Calc'd: C, 76.40; H, 4.87 Found: C,76.43; H, 5.00

- 51 -

Example 49

(2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid

5 The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol, and (R)-(+)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 90-92 °C; MS *m/e* 679 (M-H)⁺;

Analysis for: C₃₆H₂₆Br₂O₄ Calc'd: C, 63.36; H, 3.84 Found: C, 63.67; H, 3.86

10

Example 50

(2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid

15 The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3-dibromo-biphenyl-4-ol, and (R)-(+)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 80-82 °C; MS *m/e* 601 (M-H)⁺;

Analysis for: C₃₆H₂₇BrO₄ Calc'd: C, 71.65; H, 4.51 Found: C, 70.87; H, 4.55

Example 51

20 (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-methyl-pentanoic acid

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol, and (S)-(-)-hydroxyisocaproic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 81-83 °C; MS *m/e* 645 (M-H)⁺;

25 Analysis for: C₃₃H₂₈Br₂O₄ Calc'd: C, 61.13; H, 4.35 Found: C, 61.30; H, 4.06

Example 52

2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-hexanoic acid

30 The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol, and ethyl DL-2-hydroxycaproate, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 75-76 °C; MS *m/e* 645 (M-H)⁺;

Analysis for: C₃₃H₂₈Br₂O₄ Calc'd: C, 61.13; H, 4.35 Found: C, 61.27; H, 4.23

35

- 52 -

Example 53(2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-butyric acid

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol, and tert-butyl (S)-(-)-hydroxybutyrate, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 79-80 °C; MS *m/e* 617 (M-H)⁺;

Analysis for: C₃₁H₂₄Br₂O₄ Calc'd: C, 60.02; H, 3.90 Found: C, 60.41; H, 3.80

Example 5410 (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-octanoic acid

The title compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol, and tert-butyl (R)-2-hydroxyoctanoate, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 72-74°C; MS *m/e* 673 (M-H)⁺;

15 Analysis for: C₃₅H₃₂Br₂O₄ Calc'd: C, 62.15; H, 4.77 Found: C, 62.01; H, 4.71

Example 55(2S)-2-[4-(2-benzyl-benzofuran-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid

20 The title compound was prepared from 4-(2-benzyl-benzofuran-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1, steps g-h, and was obtained as a white solid, mp 73-75 °C; MS *m/e* 629 (M-H)⁺;

Analysis for: C₄₃H₃₄O₅ x 0.5 H₂O Calc'd: C, 80.73; H, 5.51 Found: C, 80.45; H, 5.32

Example 56[4-(2-benzyl-benzo[b]thiophen-3-yl)-5'-bromo-3'',4''-dimethoxy[1,1':3',1'']terphenyl]-4'-yloxy]-acetic acid

30 The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-5-bromo-3'',4''-dimethoxy-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 103-105 °C; MS *m/e* 663 (M-H)⁺;

Analysis for: C₃₇H₂₉BrO₅S Calc'd: C, 66.77; H, 4.39 Found: C, 66.77; H, 4.30

35

- 53 -

Example 57

[4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-3''-methoxy[1,1':3',1''-terphenyl]-4'-yl)oxy]-acetic acid

5 The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-5-bromo-3''-methoxy-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 88-90 °C; MS *m/e* 633 (M-H)⁺;

Analysis for: C₃₆H₂₇BrO₄S Calc'd: C, 68.03; H, 4.28 Found: C, 66.75; H, 4.18

10

Example 58

[[3,3-dimethoxy-5-[4-[2-(phenylmethyl)benzyl-benzo[b]-thiophen-3-yl]-[1,1':3',1''-terphenyl]-4'-yl)oxy]-acetic acid

15 The title compound was prepared from 3,3''-dimethoxy-5'-[4-[2-(phenylmethyl)benzo[b]thien-3-yl]phenyl][1,1':3',1''-terphenyl]-2'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 83-85 °C; MS *m/e* 661 (M-H)⁺;

Analysis for: C₄₃H₃₄O₅S x 0.8 H₂O Calc'd: C, 76.26; H, 5.30 Found: C, 76.35; H, 5.07

20

Example 59

[4-(2-benzyl-benzo[b]-thiophen-3-yl)-4''-methoxy[1,1':3',1''-terphenyl]-4'-yl)oxy]-acetic acid

25 The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 75-77 °C; MS *m/e* 555 (M-H)⁺;

Analysis for: C₃₆H₂₈O₄S x 1 H₂O Calc'd: C, 75.24; H, 5.26 Found: C, 75.56; H, 5.03

Example 60

30 [4-(2-benzyl-benzo[b]-thiophen-3-yl)-3'',4''-dimethoxy[1,1':3',1''-terphenyl]-4'-yl)oxy]-acetic acid

35 The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-3'',4''-dimethoxy-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 95-97 °C; MS *m/e* 585 (M-H)⁺;

- 54 -

Analysis for: $C_{37}H_{30}O_5S \times 0.8 H_2O$ Calc'd: C, 73.93; H, 5.30 Found: C, 73.96; H, 4.98

Example 61

5 [4-(2-benzyl-benzo[b]thiophen-3-yl)-3'',4'',5''-trimethoxy[1,1':3',1'']-terphenyl]-4'-yl)oxy]-acetic acid

The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-2'',3'',4''-trimethoxy-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained
10 as a white solid, mp 93-94 °C; MS *m/e* 615 (M-H)⁺;
Analysis for: $C_{38}H_{32}O_6S \times 1.0 H_2O$ Calc'd: C, 71.90; H, 5.40 Found: C, 72.12; H, 5.14

Example 62

15 [4-(2-benzyl-benzo[b]thiophen-3-yl)-[1,1':3',1'']-terphenyl]-4'-yl)oxy]-acetic acid

The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-[1,1':3',1'']terphenyl-4'-ol, and methyl bromoacetate, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 88-90 °C; MS *m/e* 525 (M-H)⁺;
20 Analysis for: $C_{35}H_{26}O_3S \times 0.3 H_2O$ Calc'd: C, 79.01; H, 5.04 Found: C, 79.10; H, 4.92

Example 63

4-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxymethyl]-benzoic acid

25 The title compound was prepared from 4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1':3',1'']terphenyl-4'-ol, and 4'-bromomethyl-benzoic acid methyl ester, in substantially the same manner, as described in Example 2, steps g, and was obtained as a white solid, mp 208-210 °C; MS *m/e* 509 (M-H)⁺;
Analysis for: $C_{35}H_{26}O_4$ Calc'd: C, 81.47; H, 5.20 Found: C, 81.41; H, 5.24

30

Example 64

(2R)-2-[4'-(2-Benzyl-4,5-dimethyl-thiophen-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

35 The title compound was prepared from 4'-(2-benzyl-4,5-dimethyl-thiophen-3-yl)-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same

- 55 -

manner, as described in Example 29, and was obtained as an off-white solid, mp 142-144 °C; MS *m/e* 517 (M-H)⁺;

Analysis for: C₃₄H₃₀O₃S Calc'd: C, 78.73; H, 5.83 Found: C, 78.06; H, 5.68

5

Example 65

(2R)-2-[4'-(2-Benzyl-benzofuarn-3-yl)-3,5-dimethyl-biphenyl-4-yloxy]-3-phenyl-propionic acid

The title compound was prepared from 4'-(2-benzyl-benzofuarn-3-yl)-3,5-dimethyl-biphenyl-4-ol, and (R)-(+)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps g-h, and was obtained as an off-white solid, mp 68-70 °C; MS *m/e* 551 (M-H)⁺;

Analysis for: C₃₈H₃₂O₄ Calc'd: C, 82.58; H, 5.84 Found: C, 80.67; H, 5.90

Example 66

15 (2R)-2-[4'-(2-Benzoyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

The title compound was prepared from 4'-(2-benzoyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps g-h, and was obtained as a yellow solid, MS *m/e* 584 (M-H)⁺;

Analysis for: C₃₆H₂₅NO₇ x 0.5 H₂O Calc'd: C, 72.97; H, 4.42; N, 2.36 Found: C, 72.67; H, 4.25; N, 2.39

Example 67

25 (2R)-2-[4'-(2-(Hydroxy-phenyl-methyl)-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

The title compound was prepared from 4'-[2-(hydroxy-phenyl-methyl)-benzofuarn-3-yl]-3-nitro-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps g-h, and was obtained as a brown solid, mp 121-125 °C; MS *m/e* 584 (M-H)⁺;

30 Analysis for: C₃₆H₂₇NO₇ x 0.75 H₂O Calc'd: C, 72.17; H, 4.79; N, 2.34 Found: C, 72.25; H, 4.80; N, 2.16

- 56 -

Example 68

(2R)-2-[4'-(2-Benzyl-benzofuran-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

Step a) 4'-(2-Benzyl-benzofuran-3-yl)-3-nitro-biphenyl-4-ol

- 5 Iron (III) nitrate nonahydrate (8.04 g, 19.9 mmol) was added to a solution of 4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-ol (6.8 g, 18.1 mmol) in absolute ethanol (80 mL), and the mixture was heated to 45 °C for 1.5 hour. The reaction mixture was cooled to room temperature and poured into HCl (0.1 N) and extracted with ethyl acetate 3 times. The extracts were washed with brine, dried over MgSO₄ and
- 10 concentrated *in vacuo*. Purification by flash chromatography (10% EtOAc/petroleum ether) gave the title compound as a yellow solid, mp 75 °C; MS *m/e* 420 (M - H)⁺. Analysis for C₂₇H₁₉NO₄ • 0.5 H₂O: Calcd. C, 75.34; H, 4.68; N, 3.25. Found: C, 75.6; H, 4.51; N, 3.11.

- 15 Step b) (2R)-2-[4'-(2-Benzyl-benzofuran-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid

- This compound was prepared from 4'-(2-benzyl-benzofuran-3-yl)-3-nitro-biphenyl-4-ol, and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps g-h, and was obtained as a light brown solid,
- 20 mp 77-82 °C; MS *m/e* 587 (M⁺);
Analysis for: C₃₆H₂₇NO₆ x 1 H₂O Calc'd: C, 73.59; H, 4.97; N, 2.38 Found: C, 73.89; H, 4.99; N, 2.29

Example 69

- 25 (2R)-2-[4'-(3-Benzyl-3H-imidazol[4,5-b]pyridin-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt

Step a) Benzyl-(3-nitro-pyridin-2-yl)-amine

- To a stirred solution of 2-chloro-3-nitropyridine (10 g, 63.1 mmol) in toluene (100 mL) was added in one portion benzyl amine (13.5 g, 126 mmol) and set to reflux
- 30 overnight. The reaction was cooled to room temperature and filtered. The solvent was evaporated and the residue was purified by flash chromatography (10% EtOAc/petroleum ether) affording the title compound as a yellow solid, mp 78 °C; MS *m/e* 229 (M⁺). Analysis for C₁₂H₁₁N₃O₂: Calcd. C, 62.87; H, 4.84; N, 18.33. Found: C, 63.15; H, 4.74; N, 18.28.

35

- 57 -

Step b) Benzyl-pyridine-2,3-diamine

A solution of benzyl-(3-nitro-pyridin-2-yl)-amine (5.00 g, 21.8 mmol) and tin (II) chloride dihydrate (24.6 g, 109.1 mmol) in EtOAc (100 ml) was refluxed for 2 hours. The reaction was cooled to room temperature, carefully quenched with sat. aq. NaHCO₃ (until basic), diluted with EtOAc (350 mL), stirred overnight and filtered. The biphasic filtrate was separated and the aqueous phase extracted with EtOAc. The combined organic extracts were dried (MgSO₄) and concentrated. Purification by flash chromatography (40% - 50% EtOAc/petroleum ether) gave the title compound as a light red solid, mp 88 °C; MS *m/e* 200 (M + H)⁺. Analysis for C₁₂H₁₃N₃: Calcd. C, 72.33; H, 6.58; N, 21.09. Found: C, 72.23; H, 6.68; N, 21.42.

Step c) 3-Benzyl-1,3-dihydro-imadazo[4,5-b]pyridin-2-one

Ethyl chloroformate (3.34 g, 30.7 mmol) was added to a solution of benzyl-pyridine-2,3-diamine (2.78 g, 14.0 mmol) in chloroform (70 mL), and the mixture was refluxed for 1.5 hours. The reaction mixture was washed with aqueous NaHCO₃ and water, and evaporated to dryness. The residue was subjected to column chromatography on silica gel. Elution with 50% EtOAc/chloroform gave a brown oil (a mixture of mono and bis ethoxycarbonylated compounds). A solution of this oil in absolute ethanol (15 mL) was added to a solution of sodium ethoxide (10 mmol) in absolute ethanol (15 mL), and the mixture was refluxed for 3 hours. The reaction mixture was concentrated *in vacuo*. The residue was diluted with water, neutralized with 2 N HCl and extracted with ethyl acetate. The extract was washed with water, dried over MgSO₄, and concentrated. Purification by flash chromatography (20% - 40% EtOAc/petroleum ether) gave the title compound as an orange solid, mp 172 °C; MS *m/e* 225 (M⁺). Analysis for C₁₃H₁₁N₃O: Calcd. C, 69.32; H, 4.92; N, 18.65. Found: C, 69.08; H, 5.01; N, 18.00.

Step d) 3-Benzyl-2-(4'-methoxy-biphenyl-4-yl)-3H-imadazo[4,5-b]pyridine

Phosphorus pentachloride (0.92 g, 4.44 mmol) was added to a refluxing suspension of 3-benzyl-1,3-dihydro-imadazo[4,5-b]pyridin-2-one (1.0 g, 4.44 mmol) in phosphorus oxychloride (15 mL). The mixture was refluxed for 12 hours. The solvent was then removed under reduced pressure. The residue was treated with water and basified with sodium hydroxide (5 N) with external cooling. The solution was extracted with ethyl acetate and washed with brine and dried over MgSO₄ and concentrated. The residue was purified by flash chromatography (20% - 30% EtOAc/petroleum ether) and used in the following step.

- 58 -

Step e)

The above haloimidazopyridine (0.423 g, 1.74 mmol) and tetrakis-(triphenylphosphine)palladium(0) (100 mg, 0.09 mmol) were dissolved in the minimum amount of 1,2-dimethoxyethane and stirred for 10 minutes at room temperature under a nitrogen atmosphere. 4'-Methoxy-biphenyl-4-boronic acid (0.61 g, 1.91 mmol) was added, followed by aqueous sodium carbonate (2 M, 3.5 mL). The mixture was refluxed for 12 hours, diluted with water and extracted with methylenechloride and dried over MgSO_4 and concentrated. Purification by flash chromatography (5% - 10% EtOAc/methylenechloride) gave the title compound as a white solid, mp 159 °C; MS *m/e* 391 (M^+). Analysis for $\text{C}_{26}\text{H}_{21}\text{N}_3\text{O}$ Calcd.: C, 79.77; H, 5.41; N, 10.73. Found: C, 79.48; H, 5.50; N, 10.80.

Step f) (2R)-2-[4'-(3-Benzyl-3H-imidazol[4,5-b]pyridin-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt

The title compound was prepared from 3-benzyl-2-(4'-methoxy-biphenyl-4-yl)-3H-imadazo[4,5-b]pyridine and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps d and g-h, and was obtained as a white solid, MS *m/e* 524 (M-H^+);

Analysis for: $\text{C}_{34}\text{H}_{27}\text{N}_3\text{O}_3\text{Na}$ Calc'd: C, 72.20; H, 5.15; N, 7.43 Found: C, 72.59; H, 5.02; N, 7.38

Example 70(2S)-2-[4'-(1-Benzyl-1H-benzimidazol-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acidStep a) 3-Benzyl-2-(4'-methoxy-biphenyl-4-yl)-1H- benzimidazole

This compound was prepared from 3-benzyl-2-chloro-1H- benzimidazole and 4-methoxy-biphenyl-4'-boronic acid, in substantially the same manner, as described in Example 69 step e, and was obtained as a white solid, mp 208 °C; MS *m/e* 391 ($\text{M} + \text{H}^+$).

Step b) (2S)-2-[4'-(1-Benzyl-1H-benzimidazol-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid

This compound was prepared from 1-benzyl-2-(4'-methoxy-biphenyl-4-yl)-3H-imadazo[4,5-b]pyridine and (S)-(-)-3-phenyllactic acid methyl ester, in substantially the

- 59 -

same manner, as described in Example 1 steps d and g-h, and was obtained as a light yellow solid, mp 254 °C; MS *m/e* 523 (M-H)⁺;

Analysis for: C₃₅H₂₈N₂O₃ x 0.6 H₂O Calc'd: C, 78.52; H, 5.45; N, 5.24 Found: C, 78.01; H, 5.38; N, 5.15

5

Example 71

(2S)-2-[4'-(5-acetyl-thiophen-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt

10 The title compound was prepared from 4'-(5-acetyl-thiophen-2-yl)-biphenyl-4-ol and (R)-(+)-3-phenyllactic acid methyl ester, in substantially the same manner, as described in Example 1 steps d and g-h, and was obtained as a yellow solid, MS *m/e* 443 (M+H)⁺;

Analysis for: C₂₇H₂₂O₄SNa Calc'd: C, 69.14; H, 4.62 Found: C, 69.09; H, 4.78

15 The following compounds were prepared in substantially the same manner as described in Examples 1-71

2-bromo-3-(4'-methoxy-biphenyl-4-yl)-benzofuran

20 4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-ol

2-butyl-3-(4'-methoxy-biphenyl-4-yl)-benzofuran

1-[3-(4'-methoxy-biphenyl-4-yl)-benzofuran-2-yl]-butan-1-one

25 [3-(4'-hydroxy-biphenyl-4-yl)-benzofuran-3-yl]-phenyl-methanone

4'-benzo[b]thiophen-3-yl-biphenyl-4-ol

30 3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophene

4'-(2-butyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol

4'-(2-benzyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol

35 [3-(4'-hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-phenyl-methanone

(4-fluoro-phenyl)-[3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophen-2-yl]-methanol

40 2-(4-fluoro-benzyl)-3-(4'-methoxy-biphenyl-4-yl)-benzo[b]thiophene


- 60 -

- 4'-[2-hydroxy-(4-methoxy-phenyl)-methyl]-benzo[b]thiophen-3-yl]-biphenyl-4-ol
- 1-[3-(4'-hydroxy-biphenyl-4-yl)-benzofuran-2-yl]-ethanone
- 5 4'-[2-(4-methoxy-benzyl)-benzo[b]thiophen-3-yl]-biphenyl-4-ol
- 4'-(2-ethyl-benzofuran-3-yl)-biphenyl-4-ol
- 10 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-propionic acid ethyl ester
- 4'-[2-(3,4-dimethoxy-benzyl)-benzothiophen-3-yl]-biphenyl-4-ol
- 4'-[2-(2,4-dimethoxy-benzyl)-benzothiophen-3-yl]-biphenyl-4-ol
- 15 3-(4'-methoxy-biphenyl-4-yl)-thiophen-2-carboxaldehyde
- 4-[3-(4'-hydroxy-biphenyl-4-yl)-benzo[b]thiophen-2-ylmethyl]-benzene-1,2-diol
- 20 4'-[2-(2,2-dimethyl-benzo[1,3]dioxo-5-ylmethyl)-benzothiophen-3-yl]-biphenyl-4-ol
- [3-(4-methoxy-biphenyl-4-yl)-thiophen-2-yl]-phenyl-methanol
- 2-benzyl-3-(4'-methoxy-biphenyl-4-yl)-thiophene
- 25 4'-(2-thiazole-2-ylmethyl-benzo[b]thiophen-3-yl)-biphenyl-4-ol
- 4'-[2-(4-hydroxy-benzyl)-benzo[b]thiophen-3-yl]-biphenyl-4-ol
- 30 2-benzyl-3-(4'-methoxy-biphenyl-4-yl)-furo[2,3-b]pyridine
- 4'-(2-benzyl-furo[2,3-b]pyridin-3-yl)-biphenyl-4-ol
- 4,4"-dimethoxy-5'-{4-[2-(phenylmethyl)benzo[b]thiophen-3-yl]phenyl}[1,1';3',1"-
- 35 terphenyl]-2'-ol
- 4-(2-benzyl-benzo[b]thiophen-3-yl)-5'-bromo-4"-methoxy-[1,1';3',1"]terphenyl-4'-ol
- 4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-diiodo-biphenyl-4-ol
- 40 4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-ol
- 4'-(2-benzyl-benzofuran-3-yl)-3-bromo-biphenyl-4-ol

- 61 -


WHAT IS CLAIMED IS:

1. A compound of formula I having the structure



I

5 wherein




B is carbon or nitrogen;

D is oxygen, sulfur, or nitrogen;

10 E is carbon or nitrogen;

Y is a bond, methylene, C(O), or CH(OH);

15 R^1 is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-, or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

20 R^2 and R^{2a} are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl;

25 R^3 and R^4 are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro, amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, $-NR^7(CH_2)_mCO_2H$, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3

- 62 -


heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms;

R^5 is hydrogen, alkyl of 1-6 carbon atoms, $-\text{CH}(R^8)R^9$, $-\text{C}(\text{CH}_2)_n\text{CO}_2R^{10}$,
 5 $-\text{C}(\text{CH}_3)_2\text{CO}_2R^{10}$, $-\text{CH}(R^8)(\text{CH}_2)_n\text{CO}_2R^{10}$, $-\text{CH}(R^8)\text{C}_6\text{H}_4\text{CO}_2R^{10}$.

R^6 is alkylene of 1 to 3 carbon atoms

R^7 is hydrogen or alkyl of 1 to 6 carbon atoms,

R^8 is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein
 10 the alkyl moiety contains 1-6 carbon atoms;



W is oxygen, sulfur, or nitrogen;

R^9 is $-\text{CO}_2R^{12}$, $-\text{CONHR}^{12}$, tetrazole, $-\text{PO}_3R^{12}$;

R^{10} is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of
 15 7-15 carbon atoms;

R^{12} is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;


$m = 1-3$;

$n = 1-6$;

with the proviso that when R^1 is halogen, Y is a bond;

20 or a pharmaceutically acceptable salt thereof.

2. The compound according to claim 1, wherein



wherein


25 B is carbon or nitrogen;

D is oxygen or sulfur;

or a pharmaceutically acceptable salt thereof.


- 63 -

3. The compound according to claim 1, wherein



- 5 or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 1, wherein



10

- or a pharmaceutically acceptable salt thereof.

5. The compound according to claim 1 which is (4'-benzofuran-3-yl-biphenyl-4-yloxy)-acetic acid or a pharmaceutically acceptable salt thereof.

15

6. The compound according to claim 1 which is 2-(4'-benzofuran-3-yl-biphenyl-4-yloxy)-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.

7. The compound according to claim 1 which is [4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid or a pharmaceutically acceptable salt thereof.

20

8. The compound according to claim 1 which is 2-[4'-(2-bromo-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.

25

9. The compound according to claim 1 which is 4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-ol or a pharmaceutically acceptable salt thereof.

10. The compound according to claim 1 which is [4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-acetic acid or a pharmaceutically acceptable salt thereof.

30

- 64 -

11. The compound according to claim 1 which is 2-[4'-(2-butyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 5 12. The compound according to claim 1 which is 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 10 13. The compound according to claim 1 which is 2-[4'-(2-butyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 15 14. The compound according to claim 1 which is 2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 20 15. The compound according to claim 1 which is 2-[4'-(2-benzoyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 25 16. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 30 17. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
18. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid tromethamine salt
19. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-propionic acid or a pharmaceutically acceptable salt thereof.
- 35 20. The compound according to claim 1 which is (R)-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid or a pharmaceutically acceptable salt thereof.

- 65 -

21. The compound according to claim 1 which is (2R)-2-{4'-[2-(4-fluoro-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 5 22. The compound according to claim 1 which is (2R)-2-{4'-[2-(4-methoxy-benzyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 10 23. The compound according to claim 1 which is [4'-(2-butyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-phenyl-acetic acid or a pharmaceutically acceptable salt thereof.
24. The compound according to claim 1 which is (2R)-2-{4'-[2-(hydroxy-phenyl-methyl)-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 15 25. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid methyl ester.
- 20 26. The compound according to claim 1 which is 2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-2-methyl-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
27. The compound according to claim 1 which is (2R)-2-{4'-[2-(3,4-dimethoxy-benzyl)-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 25 28. The compound according to claim 1 which is (2R)-2-{4'-[2-(2,4-dimethoxy-benzyl)-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 30 29. The compound according to claim 1 which is (2R)-2-{4'-[2-(2,2-dimethyl-benzo[1,3]dioxo-5-ylmethyl)-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 35

- 66 -

30. The compound according to claim 1 which is (2R)-2-[4'-(2-(3,4-dihydroxy-benzyl)-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 5 31. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-thiophene-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
32. The compound according to claim 1 which is 3-phenyl-2-[4'-(2-thiazole-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid or a
10 pharmaceutically acceptable salt thereof.
33. The compound according to claim 1 which is (2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid sodium salt
15
34. The compound according to claim 1 which is (2R)-3-phenyl-2-[4'-(2-pyridin-2-ylmethyl-benzo[b]thiophene-3-yl)-biphenyl-4-yloxy]-propionic acid or a pharmaceutically acceptable salt thereof.
- 20 35. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-furo[2,3]pyridin-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
36. The compound according to claim 1 which is 4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-ol or a pharmaceutically acceptable salt thereof.
25
37. The compound according to claim 1 which is 4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-ol or a pharmaceutically acceptable salt thereof.
- 30 38. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
39. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid or a
35 pharmaceutically acceptable salt thereof.

- 67 -

40. The compound according to claim 1 which is (2R)-2-[4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-methoxy-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 5 41. The compound according to claim 1 which is [(4,4''-dimethoxy-5'-{2-(phenylmethyl)benzo[b]thien-3-yl}phenyl)[1,1';3',1'']terphenyl]-2'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 10 42. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-4''-methoxy[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 15 43. The compound according to claim 1 which is [4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 20 44. The compound according to claim 1 which is [4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 25 45. The compound according to claim 1 which is (2S)-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-phenyl-butyric acid or a pharmaceutically acceptable salt thereof.
- 30 46. The compound according to claim 1 which is 4-[4'-(2-benzyl-benzo[b]thiophene-3-yl)-3-bromo-biphenyl-4-yloxy]-butyric acid or a pharmaceutically acceptable salt thereof.
- 35 47. The compound according to claim 1 which is N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-carboxy-propyl]-phthalamic acid or a pharmaceutically acceptable salt thereof.
48. The compound according to claim 1 which is N-[(3R)-3-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-methoxycarbonyl-propyl]-phthalamic acid or a pharmaceutically acceptable salt thereof.

- 68 -

49. The compound according to claim 1 which is (2R)-2-[4-(2-benzyl-benzo[b]thiophene-3-yl)-4''-chloro-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.

5

50. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-phenyl-butyric acid or a pharmaceutically acceptable salt thereof.

10

51. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzo[b]thiophen-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid or a pharmaceutically acceptable salt thereof.

15

52. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxy]-4-(1,3-dioxo-1,3-dihydro-isoindol-2-yl)-butyric acid or a pharmaceutically acceptable salt thereof.

20

53. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.

25

54. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3-bromo-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.

55. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-4-methyl-pentanoic acid or a pharmaceutically acceptable salt thereof.

30

56. The compound according to claim 1 which is 2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-hexanoic acid or a pharmaceutically acceptable salt thereof.

35

57. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-butyric acid or a pharmaceutically acceptable salt thereof.


58. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuran-3-yl)-3,5-dibromo-biphenyl-4-yloxy]-octanoic acid or a pharmaceutically acceptable salt thereof.
- 5
59. The compound according to claim 1 which is (2S)-2-[4-(2-benzyl-benzofuran-3-yl)-4''-methoxy-[1,1';3',1'']terphenyl-4'-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 10
60. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-3'',4''-dimethoxy[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 15
61. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-5'-bromo-3''-methoxy[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 20
62. The compound according to claim 1 which is [[3,3-dimethoxy-5-[4-[2-(phenylmethyl)benzyl-benzo[b]-thiophen-3-yl)-[1,1';3',1'']terphenyl]-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 25
63. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-4''-methoxy[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 30
64. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-3'',4''-dimethoxy[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
- 35
65. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-3'',4'',5''-trimethoxy[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.
66. The compound according to claim 1 which is [4-(2-benzyl-benzo[b]-thiophen-3-yl)-[1,1';3',1'']terphenyl-4'-yl)oxy]-acetic acid or a pharmaceutically acceptable salt thereof.

- 70 -

67. The compound according to claim 1 which is 4-[4'-(2-benzyl-benzofuran-3-yl)-biphenyl-4-yloxymethyl]-benzoic acid or a pharmaceutically acceptable salt thereof.
- 5 68. The compound according to claim 1 which is (2S)-2-[4'-(1-benzyl-1H-benzimidazol-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
69. The compound according to claim 1 which is (2R)-2-[4'-(3-benzyl-3H-imidazol[4,5-b]pyridin-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt.
- 10 70. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-4,5-dimethyl-thiophen-3-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 15 71. The compound according to claim 1 which is (2R)-2-[4'-(2-benzoyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 20 72. The compound according to claim 1 which is (2R)-2-[4'-(2-benzyl-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 25 73. The compound according to claim 1 which is (2R)-2-[4'-(2-(hydroxy-phenyl-methyl)-benzofuarn-3-yl)-3-nitro-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
- 30 74. The compound according to claim 1 which is (2S)-2-[4'-(2-benzyl-benzofuarn-3-yl)-3,5-dimethyl-biphenyl-4-yloxy]-3-phenyl-propionic acid or a pharmaceutically acceptable salt thereof.
75. The compound according to claim 1 which is (2S)-2-[4'-(5-acetyl-thiophen-2-yl)-biphenyl-4-yloxy]-3-phenyl-propionic acid sodium salt.


- 71 -

76. A method of treating metabolic disorders mediated by insulin resistance or hyperglycemia in a mammal in need thereof which comprises administering to said mammal, a compound of formula I having the structure



I

5 wherein




B is carbon or nitrogen;

D is oxygen, sulfur, or nitrogen;

10 E is carbon or nitrogen;

Y is a bond, methylene, C(O), or CH(OH);

15 R¹ is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-, or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

20 R² and R²ᵃ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl;

25 R³ and R⁴ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro, amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, -NR⁷(CH₂)ₘCO₂H, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3

- 72 -


heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms;

R⁵ is hydrogen, alkyl of 1-6 carbon atoms, -CH(R⁸)R⁹, -C(CH₂)_nCO₂R¹⁰,
 5 -C(CH₃)₂CO₂R¹⁰, -CH(R⁸)(CH₂)_nCO₂R¹⁰, -CH(R⁸)C₆H₄CO₂R¹⁰

R⁶ is alkylene of 1 to 3 carbon atoms

R⁷ is hydrogen or alkyl of 1 to 6 carbon atoms,

R⁸ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12
 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein
 10 the alkyl moiety contains 1-6 carbon atoms;



W is oxygen, sulfur, or nitrogen;

R⁹ is -CO₂R¹², -CONHR¹², tetrazole, -PO₃R¹²;

R¹⁰ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of
 15 7-15 carbon atoms;

R¹² is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;


m = 1-3;

n = 1-6;

with the proviso that when R¹ is halogen, Y is a bond;


20 or a pharmaceutically acceptable salt thereof.

77. A method of treating or inhibiting type II diabetes in a mammal in need thereof which comprises administering to said mammal, a compound of formula I having the structure



25 wherein

- 73 -




B is carbon or nitrogen;

D is oxygen, sulfur, or nitrogen;

5 E is carbon or nitrogen;

Y is a bond, methylene, C(O), or CH(OH);

R¹ is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-,
 10 or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

R² and R^{2a} are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-
 15 6 carbon atoms, halogen, trifluoromethyl;


R³ and R⁴ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl
 of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro,
 amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide,
 arylsulfoamide, cycloalkyl of 3-8 carbon atoms, -NR⁷(CH₂)_mCO₂H,
 20 pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3
 heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon
 atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon
 atoms or, alkoxy of 1-6 carbon atoms;

R⁵ is hydrogen, alkyl of 1-6 carbon atoms, -CH(R⁸)R⁹, -C(CH₂)_nCO₂R¹⁰,
 25 -C(CH₃)₂CO₂R¹⁰, -CH(R⁸)(CH₂)_nCO₂R¹⁰, -CH(R⁸)C₆H₄CO₂R¹⁰

R⁶ is alkylene of 1 to 3 carbon atomsR⁷ is hydrogen or alkyl of 1 to 6 carbon atoms,

- 74 -

R^8 is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein the alkyl moiety contains 1-6 carbon atoms;



5 W is oxygen, sulfur, or nitrogen;

R^9 is $-\text{CO}_2\text{R}^{12}$, $-\text{CONHR}^{12}$, tetrazole, $-\text{PO}_3\text{R}^{12}$;

R^{10} is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of 7-15 carbon atoms;

R^{12} is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;


10 $m = 1-3$;

$n = 1-6$;


with the proviso that when R^1 is halogen, Y is a bond;

or a pharmaceutically acceptable salt thereof.

15 78. A method of modulating glucose levels in a mammal in need thereof which comprises administering to said mammal, a compound of formula I having the structure



wherein



20

B is carbon or nitrogen;


D is oxygen, sulfur, or nitrogen;

E is carbon or nitrogen;

Y is a bond, methylene, $\text{C}(\text{O})$, or $\text{CH}(\text{OH})$;


- 75 -

- R^1 is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-, or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

- R^2 and R^{2a} are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl;
- R^3 and R^4 are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro, amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, $-NR^7(CH_2)_mCO_2H$, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3 heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms;
- R^5 is hydrogen, alkyl of 1-6 carbon atoms, $-CH(R^8)R^9$, $-C(CH_2)_nCO_2R^{10}$, $-C(CH_3)_2CO_2R^{10}$, $-CH(R^8)(CH_2)_nCO_2R^{10}$, $-CH(R^8)C_6H_4CO_2R^{10}$
- R^6 is alkylene of 1 to 3 carbon atoms
- R^7 is hydrogen or alkyl of 1 to 6 carbon atoms,
- R^8 is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein the alkyl moiety contains 1-6 carbon atoms;



W is oxygen, sulfur, or nitrogen;

R^9 is $-CO_2R^{12}$, $-CONHR^{12}$, tetrazole, $-PO_3R^{12}$;

- 76 -

R^{10} is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of 7-15 carbon atoms;


R^{12} is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;
 $m = 1-3$;

5 $n = 1-6$;


with the proviso that when R^1 is halogen, Y is a bond;

or a pharmaceutically acceptable salt thereof.

79. A pharmaceutical composition which comprises a compound of formula I
 10 having the structure



wherein




B is carbon or nitrogen;

15 D is oxygen, sulfur, or nitrogen;

E is carbon or nitrogen;

Y is a bond, methylene, C(O), or CH(OH);

R^1 is alkyl containing 1 to 12 carbons, aryl of 6-12 carbon atoms, arylalkyl of 7-15
 carbon atoms, halogen, carboxaldehyde, trifluoromethyl, alkoxy of 1-6 carbon
 20 atoms, 2,2-dimethyl-1,3-benzodioxole, Het-alkyl wherein the alkyl moiety
 contains 1-6 carbon atoms, or aryl of 6-10 carbon atoms which is mono-, di-,
 or tri- substituted with halogen, trifluoromethyl, or alkoxy of 1-6 carbon atoms;



G is oxygen, sulfur or nitrogen;

25

- 77 -

R² and R^{2a} are each, independently, hydrogen, alkyl of 1-6 carbon atoms, alkoxy of 1-6 carbon atoms, halogen, trifluoromethyl;


5 R³ and R⁴ are each, independently, hydrogen, halogen, alkyl of 1-6 carbon atoms, aryl of 6-10 carbon atoms, trifluoromethyl, alkoxy of 1-6 carbon atoms, nitro, amino, carboalkoxy, carbamide, carbamate, urea, alkylsulfoamide, arylsulfoamide, cycloalkyl of 3-8 carbon atoms, -NR⁷(CH₂)_mCO₂H, pyrrolidinone, a heterocyclic ring containing 5 to ring 7 atom rings having 1 to 3 heteroatoms selected from oxygen, nitrogen, or sulfur, or aryl of 6-10 carbon atoms mono-, di-, or tri-substituted with trifluoromethyl, alkyl of 1-6 carbon atoms or, alkoxy of 1-6 carbon atoms;

10 R⁵ is hydrogen, alkyl of 1-6 carbon atoms, -CH(R⁸)R⁹, -C(CH₂)_nCO₂R¹⁰, -C(CH₃)₂CO₂R¹⁰, -CH(R⁸)(CH₂)_nCO₂R¹⁰, -CH(R⁸)C₆H₄CO₂R¹⁰

R⁶ is alkylene of 1 to 3 carbon atoms

R⁷ is hydrogen or alkyl of 1 to 6 carbon atoms,

15 R⁸ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 6-12 carbon atoms, aralkyl of 6-12 carbon atoms, cycloalkyl of 3-8 carbon atoms, phthalic acid, or Q-alkyl wherein the alkyl moiety contains 1-6 carbon atoms;



W is oxygen, sulfur, or nitrogen;

20 R⁹ is -CO₂R¹², -CONHR¹², tetrazole, -PO₃R¹²;

R¹⁰ is hydrogen, alkyl of 1-6 carbon atoms, aryl of 7-15 carbon atoms, or aralkyl of 7-15 carbon atoms;

R¹² is hydrogen, alkyl, aryl of 6-12 carbon atoms, aralkyl of 7-15 carbon atoms;

m = 1-3;

25 n = 1-6;

with the proviso that when R¹ is halogen, Y is a bond;

or a pharmaceutically acceptable salt thereof, and a pharmaceutical carrier.